Math 68. Algebraic Combinatorics.

Problem Set 1. Due on Friday, 9/27/2013

1. Prove the identities

$$\sum_{i=0}^{n} \binom{i}{k} = \binom{n+1}{k+1}, \qquad \qquad \sum_{i=0}^{n} \binom{m+i}{i} = \binom{m+n+1}{n}$$

combinatorially, by showing that both sides of each equality count the same thing.

2. Give a combinatorial proof of the equality

$$\sum_{k=0}^{n} (-1)^k \binom{n}{k} = 0$$

by describing a bijection between subsets of [n] of odd size and subsets of [n] of even size.

- 3. Let $[n] = \{1, 2, \dots, n\}.$
 - (a) Find the number of k-tuples (S_1, S_2, \ldots, S_k) of subsets of [n] such that $S_1 \subseteq S_2 \subseteq \cdots \subseteq S_k$.
 - (b) Find the number of k-tuples (S_1, S_2, \ldots, S_k) of subsets of [n] such that $S_1 \cap S_2 \cap \cdots \cap S_k = \emptyset$.
- 4. A Delannoy path is a lattice path in \mathbb{Z}^2 from (0,0) to (m,n) using steps (1,0) (horizontal), (0,1) (vertical), and (1,1) (diagonal). The number of these paths is the Delannoy number $D_{m,n}$. For example, $D_{2,1} = 5$. Prove that

$$D_{m,n} = \sum_{k} \binom{m}{k} \binom{n+k}{m}.$$

Hint: Classify the paths according to the number of diagonal steps.

- 5. Prove that the number of partitions of n into odd parts equals the number of partitions of n into distinct parts.
- 6. (a) In how many ways can we choose k points, no two consecutive, from a collection of n points arranged in a line?
 - (b) What if the n points are arranged in a circle?
- 7. A set partition π of [n] is a way to subdivide [n] into nonempty blocks. A set partition is called *noncrossing* if it contains no two blocks B and B' such that $i, k \in B$ and $j, l \in B'$ for some i < j < k < l. Show that the number of noncrossing partitions equals the Catalan number $C_n = \frac{1}{n+1} {\binom{2n}{n}}$.