

# Dartmouth College

Mathematics 101

Homework 7 (due Wednesday, November 15)

## 1. Localization.

(a) Let  $A = \mathbb{Z}$  and  $\mathfrak{P} = p\mathbb{Z}$  with  $p$  a prime in  $\mathbb{Z}$ . We have characterized the localization  $A_{\mathfrak{P}} = \mathbb{Z}_{\mathfrak{P}}$  (the localization of  $\mathbb{Z}$  at the prime ideal  $\mathfrak{P}$ ) as  $\mathbb{Z}_{\mathfrak{P}} = \{a/b \in \mathbb{Q} \mid a, b \in \mathbb{Z}, p \nmid b\}$ . Show that  $\mathbb{Z}_{\mathfrak{P}}/p\mathbb{Z}_{\mathfrak{P}} \cong \mathbb{Z}/p\mathbb{Z}$ .

(b) Let  $A$  be a commutative ring with identity.

i. Suppose that for each prime ideal  $\mathfrak{P}$  in  $A$ , the local ring  $A_{\mathfrak{P}}$  has no nonzero nilpotent elements. Show that  $A$  has no nonzero nilpotent elements. *Hint:* Show that for an element  $x \in A$ , the set  $\text{Ann}(x) = \{y \in A \mid yx = 0\}$  is an ideal of  $A$ .  $\text{Ann}(x)$  is called the annihilator of the element  $x$ .

ii. Proof or counterexample: If each  $A_{\mathfrak{P}}$  is an integral domain, then  $A$  is an integral domain.

2. Let  $F$  be a field, and let  $a, b \in F^{\times}$ . Denote by  $A = \left(\frac{a, b}{F}\right)$  the quaternion algebra over  $F$  defined as follows:  $A$  is a four-dimensional vector space over  $F$  with basis  $\{1, i, j, k\}$ . The basis elements satisfy  $i^2 = a, j^2 = b, ij = k = -ji$ , and the scalars in  $F$  commute with all elements of  $A$ . In fact  $F$  is the center of  $A$ . The algebra  $\mathbb{H} = \left(\frac{-1, -1}{\mathbb{R}}\right)$  is known as Hamilton's quaternions.

(a) There is a natural involution on  $A$  denoted  $\alpha \mapsto \bar{\alpha}$  which for scalars  $w, x, y, z$  takes  $\alpha = w + xi + yj + zk$  to  $\bar{\alpha} = w - xi - yj - zk$ . Define two maps with domain  $A$  called the norm and trace, given by  $N(\alpha) = \alpha\bar{\alpha}$ , and  $Tr(\alpha) = \alpha + \bar{\alpha}$ .

i. Find explicit formulas for the norm and trace in terms of the variables  $w, x, y, z$  when  $\alpha = w + xi + yj + zk$ .

ii. Show that both the norm and trace take values in  $F$ , and prove that every element of  $A$  is the root of a quadratic equation with coefficients in  $F$ .

iii. If  $F = \mathbb{R}$ , show that  $A$  is a division ring if and only if  $a < 0$  and  $b < 0$ .

(b) Let  $R$  be a ring with identity, and let  $\alpha \in R$ . Consider the evaluation map  $\varphi_{\alpha} : R[x] \rightarrow R$  whose domain is the polynomial ring  $R[x]$ , defined by  $\varphi_{\alpha}(f) =$

$f(\alpha)$ . From class, we know that if  $R$  is commutative, then  $\varphi_\alpha$  is a ring homomorphism. Show that if  $R$  is not commutative,  $\varphi_\alpha$  is not necessarily a homomorphism. Hint: Hamilton's quaternions would be a nice ring to work with.

- (c) Consider the following popular argument in textbooks for showing a nonzero polynomial of degree  $n$  with coefficients in a field has at most  $n$  distinct roots in the field.

The proof typically proceeds by induction on  $n$ . Suppose that  $A$  is a field, and let  $f(x) \in A[x]$  have degree  $n > 1$ , and let  $\alpha \in A$  with  $f(x) = (x - \alpha)g(x)$  for  $g \in A[x]$  with degree of  $g$  equaling  $n - 1$ . Let  $\beta$  be a root of  $f$  and assume that  $\alpha \neq \beta$ . Then  $\beta$  is a root of  $g$ , and so by induction  $f$  has at most  $n$  distinct roots.

While the argument can be made rigorous in the case  $A$  is a field, it is rarely done. Given the exact argument as above, let  $A$  be a division ring (necessarily with identity). Find a counterexample to the assertion about the number of distinct roots, and explain where the gap appears in the above argument in the case of a non-commutative division ring.