

# Math 22 HW 9

- 6.1 (3 points) <sup>19</sup>
- a. True. See the definition of  $\|v\|$ .
  - b. True. See Theorem 1(c).
  - c. True. See the discussion of Fig. 5.
  - d. False. Counterexample  $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ .
  - e. True. See the box following Example 6.

6.2 (2 points) <sup>10</sup> Show  $u_1 \cdot u_2 = 0, u_2 \cdot u_3 = 0, u_3 \cdot u_1 = 0$ .

Use Theorem 4 and observe that three linearly independent vectors in  $\mathbb{R}^3$  form a basis.

$$x = \frac{4}{5}u_1 + \frac{1}{5}u_2 + \frac{1}{5}u_3$$

$\uparrow$   $\uparrow$   $\uparrow$   
 $x_1$   $x_2$   $x_3$

use projection formula  
 $x_i = \frac{x \cdot u_i}{u_i \cdot u_i} \quad i=1,2,3$

14 (2 points)  $y = \begin{bmatrix} 14/5 \\ 2/5 \end{bmatrix} + \begin{bmatrix} -4/5 \\ 29/5 \end{bmatrix}$

20 (3 points) Not orthogonal. Orthonormal set  $\begin{bmatrix} -2/3 \\ 1/3 \\ 2/3 \end{bmatrix}, \begin{bmatrix} 1/5 \\ 4/5 \\ 0 \end{bmatrix}$

means  $u_i \cdot u_j = \begin{cases} 1 & i=j \\ 0 & i \neq j \end{cases}$  ie  $U^T U = I$   
 (left alone)  $\left\{ \begin{array}{l} \text{divided by } 5^2 \end{array} \right.$

28 (2 points)  $U$  orthogonal  $\Rightarrow U U^{-1} = U U^T = I$ .

if  $V = U^T$

Theorem 6  $\Rightarrow U^T$  has orthonormal columns.

now use IMT to claim also  $U U^T = I$ , since  $U^{-1} = U^T$

then  $U U^T = I$

(In particular, the columns of  $U^T$  are linearly independent and hence form a basis for  $\mathbb{R}^n$ )

means  $V^T V = I \quad \therefore$  The rows of  $U$  form a ~~basis~~ orthonormal basis for  $\mathbb{R}^n$ .  
 ie  $V$  is orthogonal matrix.

note  
 $L.I.$   
 not relevant here

6.3 (2 points) <sup>2</sup>  $v = 2u_1 + \frac{3}{7}u_2 + \frac{12}{7}u_3 - \frac{8}{7}u_4$

8 (3 points)  $y = \begin{bmatrix} 3/2 \\ -7/2 \\ 1 \end{bmatrix} + \begin{bmatrix} -5/2 \\ 1/2 \\ 2 \end{bmatrix}$

24 (3 points) a. By hypothesis, the vectors  $w_1, \dots, w_p$  are pairwise orthogonal, and the vectors  $v_1, \dots, v_q$  are pairwise orthogonal.

Also,  $w_i \cdot v_j = 0$  for any  $i$  and  $j$  because the  $v$ 's are in the orthogonal complement of  $w$ .

b.  $y \in \mathbb{R}^n$ , write  $y = \vec{y} + z$  as in the Orthogonal Decomposition Theorem, with  $\vec{y} \in W, z \in W^\perp$

Then there exist scalars  $a_1, \dots, a_p$  and  $d_1, \dots, d_q$  such that

$$y = \vec{y} + z = a_1 w_1 + \dots + a_p w_p + d_1 v_1 + \dots + d_q v_q \quad \text{for any } \vec{y} \in \mathbb{R}^n$$

Thus  $\{w_1, \dots, w_p, v_1, \dots, v_q\}$  spans  $\mathbb{R}^n$ .

c. The set  $\{w_1, \dots, w_p, v_1, \dots, v_q\}$  is linearly independent by (a), spans  $\mathbb{R}^n$  by (b), and thus is a basis for  $\mathbb{R}^n$ . Hence  $\dim W + \dim W^\perp = p + q = \dim \mathbb{R}^n = n$ .

6.5 <sup>2</sup> (3 points) a.  $\begin{bmatrix} 12 & 8 \\ 8 & 10 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} -24 \\ -2 \end{bmatrix}$       b.  $\vec{x} = \begin{bmatrix} -4 \\ 3 \end{bmatrix}$

20 (2 points) Suppose that  $Ax = 0$ . Then  $A^T Ax = A^T 0 = 0$ .

Since  $A^T A$  is invertible by hypothesis,  $x = 0$ .

$\therefore$  The columns of  $A$  are linearly independent.

7.1 <sup>9</sup> (2 points) Orthogonal.  $U^{-1} = U^T = \begin{bmatrix} 1/\sqrt{2} & 1/\sqrt{2} \\ -1/\sqrt{2} & 1/\sqrt{2} \end{bmatrix}$  (counterclockwise rotation by  $45^\circ$ )

20 (3 points)  $P = \begin{bmatrix} 2/3 & 1/3 & 2/3 \\ 1/3 & 2/3 & 2/3 \\ 2/3 & 2/3 & 1/3 \end{bmatrix}$ ,  $D = \begin{bmatrix} 13 & 0 & 0 \\ 0 & 7 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ .

35 (3 points) a. Note that  $(u u^T)x = u(u^T x) = (u^T x)u$  because  $u^T x$  is a scalar.

$$\therefore u^T(x - Bx) = u^T x - u^T(u u^T)x = u^T x - (u^T u)u^T x = u^T x - u^T x = 0.$$

$\therefore Bx$  is the orthogonal projection of  $x$  onto  $U$ .

b.  $B_{ij} = u_i^T u_j = u_j^T u_i = B_{ji} \Rightarrow B$  is symmetric.

$$B^2 = u u^T u u^T = u(u^T u)u^T = u u^T = B$$

c.  $Bu = u u^T u = u(u^T u) = u \cdot 1 = 1 \cdot u$

$\therefore u$  is an eigen vector of  $B$  with an eigen value of 1.