

A) Is the sequence $f_n(x) = \begin{cases} 1 & x < \frac{1}{n} \\ 0 & \text{otherwise} \end{cases}$ convergent to 0, on $(0,1)$

in the sense of $\begin{cases} \text{pointwise} & ? \\ \text{uniform} & ? \\ L^2 & ? \end{cases}$ (careful: interval is $(0,1)$ not $[0,1]$)

B) Same for sequence $f_n(x) = \begin{cases} \sqrt{n} & x < \frac{1}{n} \\ 0 & \text{otherwise} \end{cases}$ on $(0,1)$.

$\begin{cases} \text{pointwise} & ? \\ \text{uniform} & ? \\ L^2 & ? \end{cases}$

C) Now consider on unbounded interval $(-\infty, \infty)$, $f_n(x) = \begin{cases} \frac{1}{n} & |x| < n \\ 0 & \text{otherwise} \end{cases}$

$\begin{cases} \text{pointwise} & ? \\ \text{uniform} & ? \\ L^2 & ? \end{cases}$


D) Modify example C) so that convergence is pointwise & uniform but not L^2 .

MATH 46 WORKSHEET :

~ SOLUTIONS ~

convergence of fumes.

Barnett
4/20/08



... etc.

A) Is the sequence

$$f_n(x) = \begin{cases} 1 & x < \frac{1}{n} \\ 0 & \text{otherwise} \end{cases}$$

sketch.

convergent to 0, on (0,1)

in the sense of

- pointwise ?
- uniform ?
- L^2 ?

yes. (for any $x \in (0,1)$, $\frac{1}{n}$ will eventually be smaller, so all f_n 's zero). (careful: interval is $(0,1)$ not $[0,1]$)

no: $\max_{x \in (0,1)} |f_n(x)| = 1 \forall n, \nrightarrow 0$

yes: $\|f_n\|^2 = \int_0^1 f_n(x)^2 dx = \frac{1}{n} \rightarrow 0$ as $n \rightarrow \infty$



it's not pointwise conv. on $[0,1]$!

Note: $n \rightarrow \infty$ plays the role of $\epsilon \rightarrow 0$ in earlier presentation of pointwise vs. uniform conv.

B) Same for sequence

$$f_n(x) = \begin{cases} \sqrt{n} & x < \frac{1}{n} \\ 0 & \text{otherwise} \end{cases}$$

- pointwise ? yes, same reason as above
- uniform ? no, since $\max_{x \in (0,1)} |f_n(x)| = \sqrt{n} \nrightarrow 0$
- L^2 ? no, since $\|f_n\|^2 = \int_0^1 f_n(x)^2 dx = \int_0^{1/n} n dx = 1 \nrightarrow 0$



Now consider on unbounded interval $(-\infty, \infty)$,

unbounded interval $(-\infty, \infty)$,

$$f_n(x) = \begin{cases} \frac{1}{n} & |x| < n \\ 0 & \text{otherwise} \end{cases}$$

- pointwise ? yes: for each $x \in \mathbb{R}$, $\lim_{n \rightarrow \infty} f_n(x) = 0$
- uniform ? yes: $\max_{x \in \mathbb{R}} |f_n(x)| = \frac{1}{n} \rightarrow 0$ as $n \rightarrow \infty$
- L^2 ? yes: $\|f_n\|^2 = \int_{-n}^n \left(\frac{1}{n}\right)^2 dx = \frac{2}{n} \rightarrow 0$ as $n \rightarrow \infty$

D) Modify example C) so that convergence is pointwise & uniform but not L^2

Change height so vanishes more slowly:

$$f_n(x) = \begin{cases} \frac{1}{n^\alpha} & |x| < n \\ 0 & \text{otherwise} \end{cases}$$

for any $\alpha \leq \frac{1}{2}$.

$$\text{then } \|f_n\|^2 = \int_{-n}^n \frac{1}{n^{2\alpha}} dx = \frac{2n^{1-2\alpha}}{1-2\alpha} \nrightarrow 0$$

Note: unif \Rightarrow pointwise always ; (ie unif. is stronger statement)

unif $\Rightarrow L^2$ on bounded interval ;

on unbounded interval unif & L^2 are indep.