

1a Evaluate: $\int 9\sqrt{x} \ln(x) dx$

We shall use integration by parts.

$$\begin{aligned} \text{Let } u &= \ln x & dv &= 9\sqrt{x} dx \\ du &= \frac{dx}{x} & v &= 9\left(\frac{2}{3}\right)x^{3/2} = 6x^{3/2} \end{aligned}$$

$$\int u dv = uv - \int v du$$

$$\begin{aligned} \int 9\sqrt{x} \ln(x) dx &= (\ln x)(6x^{3/2}) - \int (6x^{3/2}) \frac{dx}{x} \\ &= 6x^{3/2} \ln x - \int 6x^{1/2} dx \end{aligned}$$

$$= 6x^{3/2} \ln x - 6\left(\frac{2}{3}\right)x^{3/2} + C$$

$$= 6x^{3/2} \ln x - 4x^{3/2} + C$$

1 b Evaluate: $\int \frac{dx}{(4-x^2)^{3/2}}$


We shall use trig substitution.

$$x = 2 \sin \theta$$

$$dx = 2 \cos \theta d\theta \quad \text{So } \sqrt{4-x^2} = \sqrt{4-4\sin^2\theta} = 2\sqrt{1-\sin^2\theta} = 2\cos\theta$$

$$\therefore \int \frac{dx}{(4-x^2)^{3/2}} = \int \frac{2\cos\theta d\theta}{(2\cos\theta)^{3/2}} = \int \frac{d\theta}{4\cos^2\theta}$$


$$= \frac{1}{4} \int \sec^2\theta d\theta = \frac{1}{4} \tan\theta + C$$



$$\text{So, } \frac{1}{4} \tan\theta + C = \frac{1}{4} \left(\frac{x}{\sqrt{4-x^2}} \right) + C$$

NAME: _____

3. (10 points) Write down a definite integral which expresses the volume generated when the region bounded by the lines $x + y = 2$, $x = y$, and the y -axis is revolved about the x -axis.



region bded above by $y = 2 - x$
below by $y = x$

Volume generated =

$$\pi \int_0^1 (2-x)^2 dx - \pi \int_0^1 x^2 dx$$

$$= \pi \int_0^1 [(2-x)^2 - x^2] dx$$

4) Find the sum of the following series or show that it converges

$$\sum_{n=1}^{\infty} \frac{(-1)^{n-1} 2^{n-1}}{5^{n+1}}$$

This is a geometric series, we need to get it in the form $\sum_{n=0}^{\infty} ar^n$ to find the sum

$$\sum_{n=1}^{\infty} \frac{(-1)^{n-1} 2^{n-1}}{5^{n+1}} = \sum_{n=1}^{\infty} \frac{(-2)^{n-1}}{5^2 5^{n-1}} = \sum_{n=0}^{\infty} \frac{1}{25} \left(\frac{-2}{5} \right)^n = \sum_{n=0}^{\infty} \frac{1}{25} \left(\frac{-2}{5} \right)^n$$

We now have it in the proper form

and can see $a = \frac{1}{25}$, $r = \frac{-2}{5}$

clearly $|r| = \frac{2}{5} < 1$ which implies the series converges

and $\sum_{n=0}^{\infty} ar^n = \frac{a}{1-r}$

$$\text{Hence } \sum_{n=1}^{\infty} \frac{(-1)^{n-1} (2)^{n-1}}{5^{n+1}} = \sum_{n=0}^{\infty} \frac{1}{25} \left(\frac{-2}{5} \right)^n = \frac{\frac{1}{25}}{1 - \left(\frac{-2}{5} \right)} = \frac{\frac{1}{25}}{\frac{7}{5}}$$

$$= \frac{5}{7(25)} = \frac{1}{35}$$

5 Determine whether the following integral converges

$$\int_1^{\infty} \frac{2 - \sin(x)}{\sqrt{x}} dx$$

$$\frac{2 - \sin(x)}{\sqrt{x}} \geq \frac{1}{\sqrt{x}} \geq \frac{1}{x} \quad (\text{for } x \geq 1)$$

Since $\sum_{n=1}^{\infty} \frac{1}{n}$ diverges, $\sum_{n=1}^{\infty} \frac{2 - \sin(x)}{\sqrt{x}} dx$ diverges

and $\int_1^{\infty} \frac{2 - \sin(x)}{\sqrt{x}} dx$ diverges as well.

6) Determine whether the following series converges absolutely conditionally or diverges.

$$\sum_{n=1}^{\infty} \frac{(-1)^n \ln(n)}{n}$$

First we will check absolute convergence

$$\sum_{n=1}^{\infty} \left| \frac{(-1)^n \ln(n)}{n} \right| = \sum_{n=1}^{\infty} \frac{\ln n}{n}$$

now $\ln n > 1$ for $n \geq 3$

thus $\frac{\ln n}{n} > \frac{1}{n}$ for $n \geq 3$

and we know from the p-series test ($p=1$) that $\sum_{n=1}^{\infty} \frac{1}{n}$ is divergent

hence by the comparison test $\sum_{n=1}^{\infty} \frac{\ln n}{n}$ is divergent

Thus $\sum_{n=1}^{\infty} \frac{(-1)^n \ln(n)}{n}$ is not absolutely convergent.

Conditional convergence?

we will use the alternating series test
 $\sum_{n=1}^{\infty} \frac{(-1)^n \ln(n)}{n}$ is clearly alternating with $b_n = \frac{\ln n}{n}$

is it decreasing

Let $f(x) = \frac{\ln x}{x}$ if $\frac{d}{dx} f(x) < 0$ then $\frac{\ln n}{n}$ is decreasing

$$\begin{aligned} \frac{d}{dx} f(x) &= \frac{d}{dx} \left(\frac{\ln x}{x} \right) = \frac{d}{dx} (x^{-1} \ln x) = -x^{-2} \ln(x) + x^{-1} \cdot x^{-1} \\ &= -\frac{\ln(x)}{x^2} + \frac{1}{x^2} = \frac{1 - \ln(x)}{x^2} < 0 \end{aligned}$$

as long as $\ln(x) > 1$

$\ln(x) > 1$ when $x > e$ thus b_n is decreasing for $n \geq 3$.

$$\lim_{n \rightarrow \infty} b_n = 0$$

Let $f(x) = \frac{\ln x}{x}$ if $\lim_{x \rightarrow \infty} f(x) = 0$ then $\lim_{n \rightarrow \infty} b_n = 0$

now $\frac{\lim_{x \rightarrow \infty} \ln(x)}{\lim_{x \rightarrow \infty} x} = \frac{\infty}{\infty}$ is an indeterminate form

thus we can apply L'Hopital's rule

$$\lim_{x \rightarrow \infty} \frac{\ln(x)}{x} = \lim_{x \rightarrow \infty} \frac{\frac{d}{dx} \ln(x)}{\frac{d}{dx} x} = \lim_{x \rightarrow \infty} \frac{\frac{1}{x}}{1} = \lim_{x \rightarrow \infty} \frac{1}{x} = 0$$

$$\text{Thus } \lim_{n \rightarrow \infty} b_n = 0$$

Hence by the alternating series test

$$\sum_{n=1}^{\infty} \frac{(-1)^n \ln(n)}{n} \text{ converges}$$

Hence it is conditionally convergent

NAME: _____

$$7 \quad (b) \sum_{n=1}^{\infty} \cos\left(\frac{1}{n}\right)$$

$$a_n = \cos\left(\frac{1}{n}\right)$$

$$\lim_{n \rightarrow \infty} \cos\left(\frac{1}{n}\right) = \cos(0) = 1$$

Since $\lim_{n \rightarrow \infty} a_n \neq 0$, the series diverges
by
(Test for Divergence)