

# The Comparison Tests

10/06/2006

Lecture 8

# The Comparison Test

Suppose that  $\sum a_n$  and  $\sum b_n$  are series with positive terms.

1. If  $\sum b_n$  is convergent and  $a_n \leq b_n$  for all  $n$ , then  $\sum a_n$  is also convergent.
2. If  $\sum b_n$  is divergent and  $a_n \geq b_n$  for all  $n$ , then  $\sum a_n$  is also divergent.

# Examples

- $\sum_{n=1}^{\infty} \frac{5}{5n-1}$

# Examples

- $\sum_{n=1}^{\infty} \frac{5}{5n-1}$
- $\sum_{n=1}^{\infty} \frac{\sin^2(n)\sqrt{n}}{n^2}$

# Examples

- $\sum_{n=1}^{\infty} \frac{5}{5n-1}$
- $\sum_{n=1}^{\infty} \frac{\sin^2(n)\sqrt{n}}{n^2}$
- $\sum_{n=1}^{\infty} \frac{n}{n^3+2n+1}$

# Examples

- $\sum_{n=1}^{\infty} \frac{5}{5n-1}$
- $\sum_{n=1}^{\infty} \frac{\sin^2(n)\sqrt{n}}{n^2}$
- $\sum_{n=1}^{\infty} \frac{n}{n^3+2n+1}$
- $\sum_{n=1}^{\infty} \frac{1}{n!}$

# The Limit Comparison Test

Suppose that  $\sum a_n$  and  $\sum b_n$  are series with positive terms. if

$$\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = c$$

where  $c$  is a finite number and  $c > 0$ , then either both series converge or both diverge.

# Examples

Which of the following series converge, and which diverge?

- $\sum_{n=1}^{\infty} \frac{2}{3^n - 1}$

# Examples

Which of the following series converge, and which diverge?

- $\sum_{n=1}^{\infty} \frac{2}{3^n - 1}$

- $\sum_{n=1}^{\infty} \frac{2n+1}{n^2+2n+1}$

# Examples

Which of the following series converge, and which diverge?

- $\sum_{n=1}^{\infty} \frac{2}{3^n - 1}$
- $\sum_{n=1}^{\infty} \frac{2n+1}{n^2+2n+1}$
- $\sum_{n=1}^{\infty} \frac{1+n \ln n}{n^2+5}$

# Examples

Which of the following series converge, and which diverge?

- $\sum_{n=1}^{\infty} \frac{2}{3^n - 1}$

- $\sum_{n=1}^{\infty} \frac{2n+1}{n^2+2n+1}$

- $\sum_{n=1}^{\infty} \frac{1+n \ln n}{n^2+5}$

- $\sum_{n=1}^{\infty} \frac{\sin n\sqrt{n}}{4n+1}$

# Estimate Sums

- If used the Comparison Test to show that a series  $\sum a_n$  converges by comparison with a series  $\sum b_n$ , then we want to estimate the sum  $\sum a_n$  by comparing remainders.
- Consider the remainders

$$R_n = s - s_n = a_{n+1} + a_{n+2} + \dots$$

and

$$T_n = t - t_n = b_{n+1} + b_{n+2} + \dots,$$

where  $s = \sum a_n$  and  $t = \sum b_n$ .

## Example

- Use the sum of the first 100 terms to approximate the sum of the series

$$\sum_{n=1}^{\infty} \frac{1}{n^2 + 1}.$$