

# Functions of Several Variables

## Lecture 21

November 6, 2006

# Functions of two variables

- A function of two variables is a rule that assigns to each ordered pair of real numbers  $(x, y)$  in a set  $D$  a unique real number denoted by  $f(x, y)$ .

# Functions of two variables

- A function of two variables is a rule that assigns to each ordered pair of real numbers  $(x, y)$  in a set  $D$  a unique real number denoted by  $f(x, y)$ .
- The set  $D$  is the domain of  $f$  and its range is the set of values that  $f$  takes on.

# Functions of two variables

- A function of two variables is a rule that assigns to each ordered pair of real numbers  $(x, y)$  in a set  $D$  a unique real number denoted by  $f(x, y)$ .
- The set  $D$  is the domain of  $f$  and its range is the set of values that  $f$  takes on.
- We also write  $z = f(x, y)$

# Functions of two variables

- A function of two variables is a rule that assigns to each ordered pair of real numbers  $(x, y)$  in a set  $D$  a unique real number denoted by  $f(x, y)$ .
- The set  $D$  is the domain of  $f$  and its range is the set of values that  $f$  takes on.
- We also write  $z = f(x, y)$
- The variables  $x$  and  $y$  are **independent variables** and  $z$  is the **dependent variable**.

## Examples

## Examples

- Find the domain of the function

$$f(x, y) = \frac{2x + 3y}{x^2 + y^2 - 9}$$

## Examples

- Find the domain of the function

$$f(x, y) = \frac{2x + 3y}{x^2 + y^2 - 9}$$

- Find the domain and range of

$$f(x, y) = \sqrt{4 - x^2 - y^2}$$

## Definition

- If  $f$  is a function of two variables with domain  $D$ , then the graph of  $f$  is the set of all points  $(x, y, z) \in \mathbb{R}^3$  such that  $z = f(x, y)$  and  $(x, y)$  is in  $D$ .

Example

## Example

- A **linear function** is a function

$$f(x) = ax + by + c$$

## Example

- A **linear function** is a function

$$f(x) = ax + by + c$$


- The graph of such a function is a plane.

## Example

- $f(x, y) = \sin(x) + \sin(y)$

## Example

- $f(x, y) = \sin(x) + \sin(y)$




## Example

- $f(x, y) = (x^2 + y^2)e^{-x^2 - y^2}$

## Example

- $f(x, y) = (x^2 + y^2)e^{-x^2 - y^2}$



# The Cobb-Douglas production function


## Example

- $P(L, K) = bL^\alpha K^{1-\alpha}$

# The Cobb-Douglas production function

## Example

- $P(L, K) = bL^\alpha K^{1-\alpha}$
- $P(L, K) = 1.01L^{0.75} K^{0.25}$



## Definition


- The **level curves** of a function  $f$  of two variables are the curves with equations  $f(x, y) = k$ , where  $k$  is constant.

## Example

- $f(x, y) = \sin(x) + \sin(y)$


## Example

- $f(x, y) = \sin(x) + \sin(y)$



## Example


- $f(x, y) = (x^2 + y^2)e^{-x^2 - y^2}$



# The Cobb-Douglas production function

## Example

- $P(L, K) = 1.01L^{0.75}K^{0.25}$



## Definition

## Definition

- We say that a function  $f(x, y)$  has limit  $L$  as  $(x, y)$  approaches a point  $(a, b)$  and we write

$$\lim_{(x,y) \rightarrow (a,b)} f(x, y) = L$$

if we can make the values of  $f(x, y)$  as close to  $L$  as we like by taking the point  $(x, y)$  sufficiently close to the point  $(a, b)$ , but not equal to  $(a, b)$ .

## Definition

- We say that a function  $f(x, y)$  has limit  $L$  as  $(x, y)$  approaches a point  $(a, b)$  and we write

$$\lim_{(x,y) \rightarrow (a,b)} f(x, y) = L$$

if we can make the values of  $f(x, y)$  as close to  $L$  as we like by taking the point  $(x, y)$  sufficiently close to the point  $(a, b)$ , but not equal to  $(a, b)$ .

- We write also  $f(x, y) \rightarrow L$  as  $(x, y) \rightarrow (a, b)$  and

$$\lim_{x \rightarrow a, y \rightarrow b} f(x, y) = L$$



- If  $f(x, y) \rightarrow L_1$  as  $(x, y) \rightarrow (a, b)$  along a path  $C_1$  and  $f(x, y) \rightarrow L_2$  as  $(x, y) \rightarrow (a, b)$  along a path  $C_2$ , where  $L_1 \neq L_2$ , then  $\lim_{(x,y) \rightarrow (a,b)} f(x, y)$  does not exist.

- If  $f(x, y) \rightarrow L_1$  as  $(x, y) \rightarrow (a, b)$  along a path  $C_1$  and  $f(x, y) \rightarrow L_2$  as  $(x, y) \rightarrow (a, b)$  along a path  $C_2$ , where  $L_1 \neq L_2$ , then  $\lim_{(x,y) \rightarrow (a,b)} f(x, y)$  does not exist.
- **Example:** Show that

$$\lim_{(x,y) \rightarrow (0,0)} \frac{x^2 - y^2}{x^2 + y^2}$$

does not exist.

## Definition

## Definition

- A function  $f$  of two variables is called **continuous at**  $(a, b)$  if

$$\lim_{(x,y) \rightarrow (a,b)} f(x,y) = f(a,b)$$

## Definition

- A function  $f$  of two variables is called **continuous at**  $(a, b)$  if

$$\lim_{(x,y) \rightarrow (a,b)} f(x, y) = f(a, b)$$

- **Examples:** polynomials, rational, trigonometric, exponential, logarithmic functions are continuous on their domain.

## Examples

## Examples

- Find the limit

$$\lim_{(x,y) \rightarrow (0,0)} \frac{2x^2 - 4y}{\sqrt{2x^2 - 4y + 1} - 1}$$

## Examples

- Find the limit

$$\lim_{(x,y) \rightarrow (0,0)} \frac{2x^2 - 4y}{\sqrt{2x^2 - 4y + 1} - 1}$$

- Find the largest set on which the function

$$\frac{2xy}{9 - x^2 - y^2}$$

is continuous.