

# Directional Derivatives and the Gradient Vector Part 2

Lecture 25

February 28, 2007

Fact

## Fact

- *If  $f$  is a differentiable function of  $x$  and  $y$ , then  $f$  has a directional derivative in the direction of any unit vector  $\mathbf{u} = \langle a, b \rangle$  and*

$$D_{\mathbf{u}}f(x, y) = f_x(x, y)a + f_y(x, y)b.$$

## Fact

- If  $f$  is a differentiable function of  $x$  and  $y$ , then  $f$  has a directional derivative in the direction of any unit vector  $\mathbf{u} = \langle a, b \rangle$  and

$$D_{\mathbf{u}}f(x, y) = f_x(x, y)a + f_y(x, y)b.$$

- If  $f$  is a function of two variables  $x$  and  $y$ , then the **gradient** of  $f$  is the vector function  $\nabla f$  defined by

$$\nabla f(x, y) = \langle f_x(x, y), f_y(x, y) \rangle = \frac{\partial f}{\partial x} \mathbf{i} + \frac{\partial f}{\partial y} \mathbf{j}.$$

## Theorem

*Suppose that  $f$  is a differentiable function of two (or three) variables. The maximum value of the directional derivative  $D_{\mathbf{u}}f(x, y)$  is  $|\nabla f|$  and it occurs when  $\mathbf{u}$  has the same direction as the gradient vector  $\nabla f(x)$ .*

Example

## Example

- If  $f(x, y) = xe^y$ , find the rate of change of  $f$  at the point  $P(2, 0)$  in the direction from  $P$  to  $Q(\frac{1}{2}, 2)$ .

## Example

- If  $f(x, y) = xe^y$ , find the rate of change of  $f$  at the point  $P(2, 0)$  in the direction from  $P$  to  $Q(\frac{1}{2}, 2)$ .
- In what direction does  $f$  have the maximum rate of change? What is this maximum rate of change?

## Example

Suppose that the temperature at a point  $(x, y, z)$  in space is given by

$$T(x, y, z) = \frac{80}{1 + x^2 + 2y^2 + 3z^2},$$

where  $T$  is measured in degree Celsius and  $x, y, z$  in meters.

## Example

Suppose that the temperature at a point  $(x, y, z)$  in space is given by

$$T(x, y, z) = \frac{80}{1 + x^2 + 2y^2 + 3z^2},$$

where  $T$  is measured in degree Celsius and  $x, y, z$  in meters.

- In which direction does the temperature increase fastest at the point  $(1, 1, -2)$ ?

## Example

Suppose that the temperature at a point  $(x, y, z)$  in space is given by

$$T(x, y, z) = \frac{80}{1 + x^2 + 2y^2 + 3z^2},$$

where  $T$  is measured in degree Celsius and  $x, y, z$  in meters.

- In which direction does the temperature increase fastest at the point  $(1, 1, -2)$ ?
- What is the maximum rate of increase?

# Tangent Planes to Level Surfaces

## Definition

## Definition

- A level surface is a surface with equation

$$F(x, y, z) = k.$$

## Definition

- A level surface is a surface with equation

$$F(x, y, z) = k.$$

- Let  $P(x_0, y_0, z_0)$  be a point on  $S$  and let  $C$  be any curve that lies on  $S$  and passes through  $P$ .

## Definition

- A level surface is a surface with equation

$$F(x, y, z) = k.$$

- Let  $P(x_0, y_0, z_0)$  be a point on  $S$  and let  $C$  be any curve that lies on  $S$  and passes through  $P$ .
- Recall that  $C$  is described by a continuous vector function

$$\mathbf{r}(t) = \langle x(t), y(t), z(t) \rangle.$$

# Tangent Planes to Level Surfaces

Fact

## Fact

- *If  $x, y,$  and  $z$  are differentiable and  $F$  is also differentiable, we can apply the Chain Rule:*

$$\frac{\partial F}{\partial x} \frac{dx}{dt} + \frac{\partial F}{\partial y} \frac{dy}{dt} + \frac{\partial F}{\partial z} \frac{dz}{dt} = 0;$$

## Fact

- If  $x$ ,  $y$ , and  $z$  are differentiable and  $F$  is also differentiable, we can apply the Chain Rule:

$$\frac{\partial F}{\partial x} \frac{dx}{dt} + \frac{\partial F}{\partial y} \frac{dy}{dt} + \frac{\partial F}{\partial z} \frac{dz}{dt} = 0;$$

- Or

$$\nabla F \cdot \mathbf{r}'(t) = 0.$$

## Fact

- If  $x, y,$  and  $z$  are differentiable and  $F$  is also differentiable, we can apply the Chain Rule:

$$\frac{\partial F}{\partial x} \frac{dx}{dt} + \frac{\partial F}{\partial y} \frac{dy}{dt} + \frac{\partial F}{\partial z} \frac{dz}{dt} = 0;$$

- Or

$$\nabla F \cdot \mathbf{r}'(t) = 0.$$

- The gradient vector at  $P$ ,  $\nabla F(x_0, y_0, z_0)$  is perpendicular to the tangent vector  $\mathbf{r}'(t_0)$  to any curve  $C$  on  $S$  that passes through  $P$ .

# The Tangent Plane

## Definition

## Definition

- We define **the tangent plane to the level surface**  $F(x, y, z) = k$  **at**  $P(x_0, y_0, z_0)$  as the plane passes through  $P$  and has normal vector  $\nabla F(x_0, y_0, z_0)$ .

## Definition

- We define **the tangent plane to the level surface**  $F(x, y, z) = k$  at  $P(x_0, y_0, z_0)$  as the plane passes through  $P$  and has normal vector  $\nabla F(x_0, y_0, z_0)$ .
- It has equation

$$F_x(x_0, y_0, z_0)(x-x_0) + F_y(x_0, y_0, z_0)(y-y_0) + F_z(x_0, y_0, z_0)(z-z_0) = 0.$$

## Definition

## Definition

- The **normal line** to  $S$  at  $P$  is the line passing through  $P$  and perpendicular to the tangent plane.

## Definition

- The **normal line** to  $S$  at  $P$  is the line passing through  $P$  and perpendicular to the tangent plane.
- The symmetric equations are

$$\frac{x - x_0}{F_x(x_0, y_0, z_0)} = \frac{y - y_0}{F_y(x_0, y_0, z_0)} = \frac{z - z_0}{F_z(x_0, y_0, z_0)}$$

## Definition

If the equation of the surface  $S$  is of the form  $z = f(x, y)$ , that is

$$F(x, y, z) = f(x, y) - z = 0$$

then the equation of the tangent plane becomes

$$f_x(x_0, y_0)(x - x_0) + f_y(x_0, y_0)(y - y_0) - (z - z_0) = 0.$$

## Examples

## Examples

- Find the tangent plane and normal line of the surface

$$F(x, y, z) = x^2 + y^2 + z - 9 = 0$$

at the point  $P_0(1, 2, 4)$ .

## Examples

- Find the tangent plane and normal line of the surface

$$F(x, y, z) = x^2 + y^2 + z - 9 = 0$$

at the point  $P_0(1, 2, 4)$ .

- Find the equation of the tangent plane at the point  $(-2, 1, -3)$  to the ellipsoid

$$\frac{x^2}{4} + y^2 + \frac{z^2}{9} = 3.$$

# Significance of the Gradient Vector

Fact

## Fact

- *The gradient  $\nabla f$  gives the direction of fastest increase of  $f$ .*

## Fact

- The gradient  $\nabla f$  gives the direction of fastest increase of  $f$ .
- The gradient  $\Delta f$  is orthogonal to the level surface  $S$  of  $f$  through a point  $P$ .