

# Grid Pattern Classes

Nik Ruškuc

`nik@mcs.st-and.ac.uk`

School of Mathematics and Statistics, University of St Andrews

Permutation Patterns, Dartmouth College, 11 Aug 2010



University  
of  
St Andrews

# Gridding a permutation




## Gridding a permutation

3, 5, 4, 6, 9, 2, 11, 12, 1, 10, 8, 7




## Gridding a permutation




# Gridding a permutation

3, 5, 4, 6, 9, 2, 11, 12, 1, 10, 8, 7



## Gridding a permutation

3, 5, 4, 6, 9, 2, 11, 12, 1, 10, 8, 7



$$M = \begin{pmatrix} 0 & 1 & -1 \\ 1 & -1 & 0 \end{pmatrix}$$

# Basic definitions



# Basic definitions

- ▶ **Gridding matrix:**  $M = (m_{ij})_{p \times q}$ ,  $m_{ij} \in \{0, \pm 1\}$ .



## Basic definitions

- ▶ Gridding matrix:  $M = (m_{ij})_{p \times q}$ ,  $m_{ij} \in \{0, \pm 1\}$ .
- ▶ Permutation  $\pi$  **admits an  $M$ -gridding** if the  $xy$ -plane with the graph  $\Gamma$  of  $\pi$  plotted in it can be partitioned into an axis parallel rectangular grid  $C_{ij}$  ( $i \in [p]$ ,  $j \in [q]$ ) such that

$$\Gamma \cap C_{ij} \text{ is } \begin{cases} \text{increasing,} & \text{if } m_{ij} = 1 \\ \text{decreasing,} & \text{if } m_{ij} = -1 \\ \emptyset, & \text{if } m_{ij} = 0. \end{cases}$$

## Basic definitions

- ▶ Gridding matrix:  $M = (m_{ij})_{p \times q}$ ,  $m_{ij} \in \{0, \pm 1\}$ .
- ▶ Permutation  $\pi$  admits an  $M$ -gridding if the  $xy$ -plane with the graph  $\Gamma$  of  $\pi$  plotted in it can be partitioned into an axis parallel rectangular grid  $C_{ij}$  ( $i \in [p]$ ,  $j \in [q]$ ) such that

$$\Gamma \cap C_{ij} \text{ is } \begin{cases} \text{increasing,} & \text{if } m_{ij} = 1 \\ \text{decreasing,} & \text{if } m_{ij} = -1 \\ \emptyset, & \text{if } m_{ij} = 0. \end{cases}$$

- ▶ The (pattern class) of all  $\pi$  that admit an  $M$ -gridding is called a **(monotone) grid class**, and denoted  $\text{Grid}(M)$ .

# Outline & Credits

- ▶ Early occurrences of grid classes in literature.
- ▶ New project: Basic properties of grid classes.
- ▶ Interplay of permutation, geometrical and language-theoretical methods.
- ▶ Future directions.

Collaborators: Michael Albert (Otago), Mike Atkinson (Otago), Mathilde Bouvel (Paris/Bordeaux), Robert Brignall (Bristol/OU), Vincent Vatter (Dartmouth/Florida), NR.



# Easy examples



# Easy examples

Example

Matrix:  $( 1 \ 1 )$



# Easy examples

## Example

Matrix:  $( 1 \ 1 )$

Description: juxtapositions of two increasing sequences.



# Easy examples

## Example

Matrix:  $\begin{pmatrix} 1 & 1 \end{pmatrix}$

Description: juxtapositions of two increasing sequences.

Basis: 321, 3142, 2143.



# Easy examples

## Example

Matrix:  $\begin{pmatrix} 1 & 1 \end{pmatrix}$

Description: juxtapositions of two increasing sequences.

Basis: 321, 3142, 2143.

Enumeration:  $2^n - n$ .



# Easy examples

## Example

Matrix:  $\begin{pmatrix} 1 & 1 \end{pmatrix}$

Description: juxtapositions of two increasing sequences.

Basis: 321, 3142, 2143.

Enumeration:  $2^n - n$ .

## Example

Matrix:  $\begin{pmatrix} -1 & 1 \end{pmatrix}$

Description: juxtapositions of a decreasing and an increasing sequence.

Basis: 231, 132.

Enumeration:  $2^{n-1}$ .



## Further examples (1)

### Example

Atkinson (1999) proves

$$\text{Av}(321, 2143) = \text{Grid} \left( \begin{array}{cc} 1 & 1 \end{array} \right) \cup \text{Grid} \left( \begin{array}{c} 1 \\ 1 \end{array} \right)$$

and derives as a consequence the enumeration for  $\text{Av}(321, 2143)$ :

$$2^{n+1} - 2n - 1 - \binom{n+1}{3}.$$



## Further examples (2)

### Example

Murphy (2003) expresses several classes with two basis permutations of lengths 3 and 4 as grid classes. For instance

$$\text{Av}(132, 4312) = \text{Grid} \left( \begin{array}{ccc} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{array} \right).$$

## Other occurrences in literature

- ▶ Atkinson (1998) – skew merge permutations.
- ▶ Profile classes (Atkinson (1999)) – permutation matrices.
- ▶  $W$ -classes (Atkinson, Murphy, NR (2002)) –  $1 \times q$  matrices.
- ▶ Murphy, Vatter (2002) – PWO.
- ▶ Waton (2007) – atomicity, Grid  $\begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$ .
- ▶ Huczynska, Vatter (2006) – sparse matrices; decidability for subclasses of grid classes.
- ▶ Vatter (submitted) – small growth rates.
- ▶ Brendan 25,081: Blend includes art arranged in rectangular array.



## Graph $\Gamma(M)$ : definition

$M = (m_{ij})_{p \times q}$  – gridding matrix.

Define graph  $\Gamma = \Gamma(M)$  as follows:



## Graph $\Gamma(M)$ : definition

$M = (m_{ij})_{p \times q}$  – gridding matrix.

Define graph  $\Gamma = \Gamma(M)$  as follows:

vertices:  $r_1, \dots, r_p, c_1, \dots, c_q$ .



## Graph $\Gamma(M)$ : definition

$M = (m_{ij})_{p \times q}$  – gridding matrix.

Define graph  $\Gamma = \Gamma(M)$  as follows:

vertices:  $r_1, \dots, r_p, c_1, \dots, c_q$ .

edges:  $r_i \sim c_j \Leftrightarrow m_{ij} = \pm 1$ .



## Graph $\Gamma(M)$ : definition

$M = (m_{ij})_{p \times q}$  – gridding matrix.

Define graph  $\Gamma = \Gamma(M)$  as follows:

vertices:  $r_1, \dots, r_p, c_1, \dots, c_q$ .

edges:  $r_i \sim c_j \Leftrightarrow m_{ij} = \pm 1$ .

### Definition

If  $\Gamma(M)$  is a forest we say that  $\text{Grid}(M)$  is a **forest grid class**.




# Graph $\Gamma(M)$ : examples

## Example

	<b>1</b>	<b>-1</b>
<b>1</b>	<b>-1</b>	


# Graph $\Gamma(M)$ : examples

## Example




# Graph $\Gamma(M)$ : examples

## Example




# Graph $\Gamma(M)$ : examples

## Example




# Graph $\Gamma(M)$ : examples

## Example




# Graph $\Gamma(M)$ : examples

## Example



# Graph $\Gamma(M)$ : examples

## Example



# Partial well order

## Definition

A pattern class is **partially well ordered (PWO)** if it doesn't contain an infinite antichain.



# Partial well order

## Definition

A pattern class is partially well ordered (PWO) if it doesn't contain an infinite antichain.

## Proposition

*A finitely based class is PWO iff it has  $\leq \aleph_0$  subclasses iff all its subclasses are finitely based.*



# Partial well order

## Definition

A pattern class is partially well ordered (PWO) if it doesn't contain an infinite antichain.

## Proposition

*A finitely based class is PWO iff it has  $\leq \aleph_0$  subclasses iff all its subclasses are finitely based.*

## Theorem (Murphy, Vatter (2002))

*A grid class is PWO iff it is a forest grid class.*



## Important things

*He took the biscuits carefully out of the packet and laid them face upward on the grass, in order as he felt of edibility. They were the same as always, a Ginger, an Osborne, a Digestive, a Petit Beurre and one anonymous. He always ate the first-named last, because he liked it the best, and the anonymous first, because he thought it very likely the least palatable. The order in which he ate the remaining three was indifferent to him and varied irregularly from day to day. On his knees now before the 5 it struck him for the first time that this reduced to a paltry six the number of ways in which he could make his meal. (...) Even if he conquered his prejudice against the anonymous, still there would be only twenty-four ways in which the biscuits could be eaten. But were he to take the final step and overcome his infatuation with the ginger, then the assortment would spring to life before him, dancing the radiant measure of its total permutability, edible in a hundred and twenty ways!*

*(S. Beckett, Murphy)*



# Geometric classes

- ▶ To every finite set  $F$  of points in a  $xy$ -plane which are  $xy$ -independent there corresponds a unique permutation  $\pi(F)$ .



## Geometric classes

- ▶ To every finite set  $F$  of points in a  $xy$ -plane which are  $xy$ -independent there corresponds a unique permutation  $\pi(F)$ .
- ▶ Let  $\mathcal{F}$  be any figure (set of points) in a plane. Define

$$\text{Sub}(\mathcal{F}) = \{\pi(F) : F \subseteq \mathcal{F}, |F| < \infty\}.$$



## Geometric classes

- ▶ To every finite set  $F$  of points in a  $xy$ -plane which are  $xy$ -independent there corresponds a unique permutation  $\pi(F)$ .
- ▶ Let  $\mathcal{F}$  be any figure (set of points) in a plane. Define

$$\text{Sub}(\mathcal{F}) = \{\pi(F) : F \subseteq \mathcal{F}, |F| < \infty\}.$$

- ▶ This is clearly a pattern class.



## Geometric classes

- ▶ To every finite set  $F$  of points in a  $xy$ -plane which are  $xy$ -independent there corresponds a unique permutation  $\pi(F)$ .
- ▶ Let  $\mathcal{F}$  be any figure (set of points) in a plane. Define

$$\text{Sub}(\mathcal{F}) = \{\pi(F) : F \subseteq \mathcal{F}, |F| < \infty\}.$$

- ▶ This is clearly a pattern class.
- ▶ Waton (et al.): permutations from parallel lines; circle permutations; convex permutations; etc.



## Geometric grid classes

To every grid class  $\text{Grid}(M)$  associate a **geometric grid class**  $\text{GGrid}(M)$  as follows:



## Geometric grid classes

To every grid class  $\text{Grid}(M)$  associate a **geometric grid class**  $\text{GGrid}(M)$  as follows:

- ▶ Draw a  $p \times q$  rectangular grid with cells  $C_{ij}$ .

## Geometric grid classes

To every grid class  $\text{Grid}(M)$  associate a **geometric grid class**  $\text{GGrid}(M)$  as follows:

- ▶ Draw a  $p \times q$  rectangular grid with cells  $C_{ij}$ .
- ▶ If  $m_{ij} = 1$  draw a SW-NE diagonal in  $C_{ij}$ .



## Geometric grid classes

To every grid class  $\text{Grid}(M)$  associate a **geometric grid class**  $\text{GGrid}(M)$  as follows:

- ▶ Draw a  $p \times q$  rectangular grid with cells  $C_{ij}$ .
- ▶ If  $m_{ij} = 1$  draw a SW-NE diagonal in  $C_{ij}$ .
- ▶ If  $m_{ij} = -1$  draw a NW-SE diagonal in  $C_{ij}$ .



## Geometric grid classes

To every grid class  $\text{Grid}(M)$  associate a **geometric grid class**  $\text{GGrid}(M)$  as follows:

- ▶ Draw a  $p \times q$  rectangular grid with cells  $C_{ij}$ .
- ▶ If  $m_{ij} = 1$  draw a SW-NE diagonal in  $C_{ij}$ .
- ▶ If  $m_{ij} = -1$  draw a NW-SE diagonal in  $C_{ij}$ .
- ▶ Let  $\mathcal{F}$  be the figure consisting of these diagonals and  $\text{GGrid}(M) = \text{Sub}(\mathcal{F})$ .



## Geometric grid classes

To every grid class  $\text{Grid}(M)$  associate a **geometric grid class**  $\text{GGrid}(M)$  as follows:


- ▶ Draw a  $p \times q$  rectangular grid with cells  $C_{ij}$ .
- ▶ If  $m_{ij} = 1$  draw a SW-NE diagonal in  $C_{ij}$ .
- ▶ If  $m_{ij} = -1$  draw a NW-SE diagonal in  $C_{ij}$ .
- ▶ Let  $\mathcal{F}$  be the figure consisting of these diagonals and  $\text{GGrid}(M) = \text{Sub}(\mathcal{F})$ .

	1	-1
1	-1	

## Geometric grid classes

To every grid class  $\text{Grid}(M)$  associate a **geometric grid class**  $\text{GGrid}(M)$  as follows:


- ▶ Draw a  $p \times q$  rectangular grid with cells  $C_{ij}$ .
- ▶ If  $m_{ij} = 1$  draw a SW-NE diagonal in  $C_{ij}$ .
- ▶ If  $m_{ij} = -1$  draw a NW-SE diagonal in  $C_{ij}$ .
- ▶ Let  $\mathcal{F}$  be the figure consisting of these diagonals and  $\text{GGrid}(M) = \text{Sub}(\mathcal{F})$ .



## Geometric grid classes

To every grid class  $\text{Grid}(M)$  associate a **geometric grid class**  $\text{GGrid}(M)$  as follows:

- ▶ Draw a  $p \times q$  rectangular grid with cells  $C_{ij}$ .
- ▶ If  $m_{ij} = 1$  draw a SW-NE diagonal in  $C_{ij}$ .
- ▶ If  $m_{ij} = -1$  draw a NW-SE diagonal in  $C_{ij}$ .
- ▶ Let  $\mathcal{F}$  be the figure consisting of these diagonals and  $\text{GGrid}(M) = \text{Sub}(\mathcal{F})$ .



## Geometric grid classes

To every grid class  $\text{Grid}(M)$  associate a **geometric grid class**  $\text{GGrid}(M)$  as follows:

- ▶ Draw a  $p \times q$  rectangular grid with cells  $C_{ij}$ .
- ▶ If  $m_{ij} = 1$  draw a SW-NE diagonal in  $C_{ij}$ .
- ▶ If  $m_{ij} = -1$  draw a NW-SE diagonal in  $C_{ij}$ .
- ▶ Let  $\mathcal{F}$  be the figure consisting of these diagonals and  $\text{GGrid}(M) = \text{Sub}(\mathcal{F})$ .

### Proposition

- ▶  $\text{GGrid}(M) \subseteq \text{Grid}(M)$ .



## Geometric grid classes

To every grid class  $\text{Grid}(M)$  associate a **geometric grid class**  $\text{GGrid}(M)$  as follows:

- ▶ Draw a  $p \times q$  rectangular grid with cells  $C_{ij}$ .
- ▶ If  $m_{ij} = 1$  draw a SW-NE diagonal in  $C_{ij}$ .
- ▶ If  $m_{ij} = -1$  draw a NW-SE diagonal in  $C_{ij}$ .
- ▶ Let  $\mathcal{F}$  be the figure consisting of these diagonals and  $\text{GGrid}(M) = \text{Sub}(\mathcal{F})$ .

### Proposition

- ▶  $\text{GGrid}(M) \subseteq \text{Grid}(M)$ .
- ▶  $\text{GGrid}(M) = \text{Grid}(M)$  if  $\Gamma(M)$  is a forest.



# Finite basis (in words)

## Theorem

*Every forest grid class  $\text{Grid}(M)$  is finitely based.*



# Finite basis (in words)

## Theorem

*Every forest grid class  $\text{Grid}(M)$  is finitely based.*

## Proof

- ▶ Every basis permutation is a one point extension of a permutation from  $\text{Grid}(M)$ .



# Finite basis (in words)

## Theorem

*Every forest grid class  $\text{Grid}(M)$  is finitely based.*

## Proof

- ▶ Every basis permutation is a one point extension of a permutation from  $\text{Grid}(M)$ .
- ▶ There are only finitely many ways to insert a point into the geometrical representation of  $\text{Grid}(M)$ .



# Finite basis (in words)

## Theorem

*Every forest grid class  $\text{Grid}(M)$  is finitely based.*

## Proof

- ▶ Every basis permutation is a one point extension of a permutation from  $\text{Grid}(M)$ .
- ▶ There are only finitely many ways to insert a point into the geometrical representation of  $\text{Grid}(M)$ .
- ▶ Re-grid: every basis element also belongs to a forest grid class.



# Finite basis (in words)

## Theorem

*Every forest grid class  $\text{Grid}(M)$  is finitely based.*

## Proof

- ▶ Every basis permutation is a one point extension of a permutation from  $\text{Grid}(M)$ .
- ▶ There are only finitely many ways to insert a point into the geometrical representation of  $\text{Grid}(M)$ .
- ▶ Re-grid: every basis element also belongs to a forest grid class.
- ▶ Use PWO.

# Finite basis (in words)

## Theorem

*Every forest grid class  $\text{Grid}(M)$  is finitely based.*

## Proof

- ▶ Every basis permutation is a one point extension of a permutation from  $\text{Grid}(M)$ .
- ▶ There are only finitely many ways to insert a point into the geometrical representation of  $\text{Grid}(M)$ .
- ▶ Re-grid: every basis element also belongs to a forest grid class.
- ▶ Use PWO.

## Theorem

*Every subclass of a forest grid class is finitely based.*




# Finite basis (in pictures)

## Theorem

*Every forest grid class  $\text{Grid}(M)$  is finitely based.*

## Proof




# Finite basis (in pictures)

## Theorem

*Every forest grid class  $\text{Grid}(M)$  is finitely based.*

## Proof




# Finite basis (in pictures)

## Theorem

*Every forest grid class  $\text{Grid}(M)$  is finitely based.*

## Proof




# Finite basis (in pictures)

## Theorem

*Every forest grid class  $\text{Grid}(M)$  is finitely based.*

## Proof




# Finite basis (in pictures)

## Theorem

*Every forest grid class  $\text{Grid}(M)$  is finitely based.*

## Proof




# Finite basis (in pictures)

## Theorem

*Every forest grid class  $\text{Grid}(M)$  is finitely based.*

## Proof




# Finite basis (in pictures)

## Theorem

*Every forest grid class  $\text{Grid}(M)$  is finitely based.*

## Proof




# Finite basis (in pictures)

## Theorem

*Every forest grid class  $\text{Grid}(M)$  is finitely based.*

## Proof




# Finite basis (in pictures)

## Theorem

*Every forest grid class  $\text{Grid}(M)$  is finitely based.*

## Proof






# Finite basis (in pictures)

## Theorem

*Every forest grid class  $\text{Grid}(M)$  is finitely based.*

## Proof



# Finite basis (in pictures)

## Theorem

Every forest grid class  $\text{Grid}(M)$  is finitely based.

## Proof

0	0	0	0	0	0	0	0	1	-1	0	0	0	0
0	0	0	0	0	0	0	0	0	0	0	0	0	0
0	0	0	0	0	0	1	0	0	0	0	-1	0	0
0	0	0	0	0	0	0	0	0	0	0	0	0	0
0	0	0	0	0	1	0	0	0	0	0	0	0	-1
0	0	0	0	1	-1	0	0	0	0	0	0	0	0
0	0	0	0	0	0	0	1	0	0	0	0	0	0
0	0	1	0	0	0	0	-1	0	0	0	0	0	0
0	0	0	0	0	0	0	0	0	0	0	0	0	0
1	0	0	0	0	0	0	0	-1	0	0	0	0	0

# Finite basis (in pictures)

## Theorem

*Every forest grid class  $\text{Grid}(M)$  is finitely based.*

## Proof

0	0	0	0	0	0	0	0	0	1	-1	0	0	0	0
0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
0	0	0	0	0	0	0	1	0	0	0	0	-1	0	0
0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
0	0	0	0	0	1	0	0	0	0	0	0	0	0	-1
0	0	0	0	1	-1	0	0	0	0	0	0	0	0	0
0	0	0	0	0	0	0	1	0	0	0	0	0	0	0
0	0	1	0	0	0	0	-1	0	0	0	0	0	0	0
0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
1	0	0	0	0	0	0	0	0	-1	0	0	0	0	0

## Corollary

*Every subclass of a forest grid class is finitely based.*

# Words are good



# Words are good

- ▶ A – an alphabet.



# Words are good

- ▶  $A$  – an alphabet.
- ▶  $A^*$  – the free monoid.



# Words are good

- ▶  $A$  – an alphabet.
- ▶  $A^*$  – the free monoid.
- ▶  $\preceq$  – the subsequence (subword) ordering.



# Words are good

- ▶  $A$  – an alphabet.
- ▶  $A^*$  – the free monoid.
- ▶  $\preceq$  – the subsequence (subword) ordering.
- ▶  $(A^*, \preceq)$  is PWO (Higman).
- ▶ All closed subsets of  $A^*$  are finitely based.



# Words are good

- ▶  $A$  – an alphabet.
- ▶  $A^*$  – the free monoid.
- ▶  $\preceq$  – the subsequence (subword) ordering.
- ▶  $(A^*, \preceq)$  is PWO (Higman).
- ▶ All closed subsets of  $A^*$  are finitely based.
- ▶ Every closed subset of  $A^*$  is a finite union of sets of the form

$$a_1 A_1^* a_2 A_2^* \dots a_k A_k^* a_{k+1}$$

with  $a_i \in A$ ,  $A_j \subseteq A$ .



# Words are good

- ▶  $A$  – an alphabet.
- ▶  $A^*$  – the free monoid.
- ▶  $\preceq$  – the subsequence (subword) ordering.
- ▶  $(A^*, \preceq)$  is PWO (Higman).
- ▶ All closed subsets of  $A^*$  are finitely based.
- ▶ Every closed subset of  $A^*$  is a finite union of sets of the form

$$a_1 A_1^* a_2 A_2^* \dots a_k A_k^* a_{k+1}$$

with  $a_i \in A$ ,  $A_j \subseteq A$ .

- ▶ Every closed subset of  $A^*$  has a rational generating function.



## Encoding gridded permutations as words

# Decoding words

- ▶ The 'inverse' process.
- ▶  $M = (m_{ij})_{p \times q}$ ,  $\Gamma(M)$  a forest.
- ▶  $A = \{a_{ij} : i \in [p], j \in [q]\}$ .
- ▶  $\phi : A^* \rightarrow \text{Grid}(M)$ .



# Decoding words

- ▶ The 'inverse' process.
- ▶  $M = (m_{ij})_{p \times q}$ ,  $\Gamma(M)$  a forest.
- ▶  $A = \{a_{ij} : i \in [p], j \in [q]\}$ .
- ▶  $\phi : A^* \rightarrow \text{Grid}(M)$ .
- ▶  $\phi$  is onto



# Decoding words

- ▶ The 'inverse' process.
- ▶  $M = (m_{ij})_{p \times q}$ ,  $\Gamma(M)$  a forest.
- ▶  $A = \{a_{ij} : i \in [p], j \in [q]\}$ .
- ▶  $\phi : A^* \rightarrow \text{Grid}(M)$ .
- ▶  $\phi$  is onto, not 1-1.



# Decoding words

- ▶ The 'inverse' process.
- ▶  $M = (m_{ij})_{p \times q}$ ,  $\Gamma(M)$  a forest.
- ▶  $A = \{a_{ij} : i \in [p], j \in [q]\}$ .
- ▶  $\phi : A^* \rightarrow \text{Grid}(M)$ .
- ▶  $\phi$  is onto, not 1-1.
- ▶  $\phi$  is length preserving,



# Decoding words

- ▶ The 'inverse' process.
- ▶  $M = (m_{ij})_{p \times q}$ ,  $\Gamma(M)$  a forest.
- ▶  $A = \{a_{ij} : i \in [p], j \in [q]\}$ .
- ▶  $\phi : A^* \rightarrow \text{Grid}(M)$ .
- ▶  $\phi$  is onto, not 1-1.
- ▶  $\phi$  is length preserving, finite-to-one.



# Decoding words

- ▶ The 'inverse' process.
- ▶  $M = (m_{ij})_{p \times q}$ ,  $\Gamma(M)$  a forest.
- ▶  $A = \{a_{ij} : i \in [p], j \in [q]\}$ .
- ▶  $\phi : A^* \rightarrow \text{Grid}(M)$ .
- ▶  $\phi$  is onto, not 1-1.
- ▶  $\phi$  is length preserving, finite-to-one.
- ▶  $w_1 \preceq w_2 \Rightarrow \phi(w_1) \preceq \phi(w_2)$ .



# Immediate corollaries

## Theorem

*Every forest grid class is PWO.*



# Immediate corollaries

## Theorem

*Every forest grid class is PWO.*

## Theorem

*Every subclass of a forest grid class is a finite union of forest grid classes.*



# Immediate corollaries

## Theorem

*Every forest grid class is PWO.*

## Theorem

*Every subclass of a forest grid class is a finite union of forest grid classes.*

## Remark

Here 'forest grid class' stands for a small generalisation: one point cells are also allowed.



## Ambiguities


If we could find a 'nice' subset  $W \subseteq A^*$  such that  $\phi \upharpoonright_W$  is a bijection between  $W$  and  $\text{Grid}(M)$  we would get a rational generating function for  $\text{Grid}(M)$ .



# Ambiguities

If we could find a 'nice' subset  $W \subseteq A^*$  such that  $\phi \upharpoonright_W$  is a bijection between  $W$  and  $\text{Grid}(M)$  we would get a rational generating function for  $\text{Grid}(M)$ .

The following make  $\phi$  non-injective:




# Ambiguities

If we could find a 'nice' subset  $W \subseteq A^*$  such that  $\phi \upharpoonright_W$  is a bijection between  $W$  and  $\text{Grid}(M)$  we would get a rational generating function for  $\text{Grid}(M)$ .

The following make  $\phi$  non-injective:

- ▶  $xy$ -independent cells can be read off in either order.




# Ambiguities

If we could find a 'nice' subset  $W \subseteq A^*$  such that  $\phi \upharpoonright_W$  is a bijection between  $W$  and  $\text{Grid}(M)$  we would get a rational generating function for  $\text{Grid}(M)$ .

The following make  $\phi$  non-injective:

- ▶  $xy$ -independent cells can be read off in either order.




# Ambiguities

If we could find a 'nice' subset  $W \subseteq A^*$  such that  $\phi \upharpoonright_W$  is a bijection between  $W$  and  $\text{Grid}(M)$  we would get a rational generating function for  $\text{Grid}(M)$ .

The following make  $\phi$  non-injective:

- ▶  $xy$ -independent cells can be read off in either order.




# Ambiguities

If we could find a 'nice' subset  $W \subseteq A^*$  such that  $\phi \upharpoonright_W$  is a bijection between  $W$  and  $\text{Grid}(M)$  we would get a rational generating function for  $\text{Grid}(M)$ .

The following make  $\phi$  non-injective:

- ▶  $xy$ -independent cells can be read off in either order.




# Ambiguities

If we could find a 'nice' subset  $W \subseteq A^*$  such that  $\phi \upharpoonright_W$  is a bijection between  $W$  and  $\text{Grid}(M)$  we would get a rational generating function for  $\text{Grid}(M)$ .

The following make  $\phi$  non-injective:

- ▶  $xy$ -independent cells can be read off in either order.
- ▶ Multiple  $M$ -griddings.




# Ambiguities

If we could find a 'nice' subset  $W \subseteq A^*$  such that  $\phi \upharpoonright_W$  is a bijection between  $W$  and  $\text{Grid}(M)$  we would get a rational generating function for  $\text{Grid}(M)$ .

The following make  $\phi$  non-injective:

- ▶  $xy$ -independent cells can be read off in either order.
- ▶ Multiple  $M$ -griddings.




# Ambiguities

If we could find a 'nice' subset  $W \subseteq A^*$  such that  $\phi \upharpoonright_W$  is a bijection between  $W$  and  $\text{Grid}(M)$  we would get a rational generating function for  $\text{Grid}(M)$ .

The following make  $\phi$  non-injective:

- ▶  $xy$ -independent cells can be read off in either order.
- ▶ Multiple  $M$ -griddings.



# Resolving ambiguities

Independent cells:

- ▶  $a_{ij}a_{st} = a_{st}a_{ij}$  where  $i \neq s, j \neq t$ .
- ▶ Trace monoids, regular sets of representatives with uniqueness.

Multiple griddings:

- ▶ There are only finitely many different gridline movements, relative to the geometric representation.
- ▶ An argument similar to basis and subclasses.

## Corollary

*Every forest grid class has a rational generating function.*



# Summary of results

For a forest grid class  $\text{Grid}(M)$  the following hold:

- ▶  $\text{Grid}(\mathcal{M})$  is finitely based.
- ▶  $\text{Grid}(\mathcal{M})$  is partially well ordered.
- ▶ Every subclass of  $\text{Grid}(\mathcal{M})$  is a finite union of (slightly generalised) forest grid classes.
- ▶  $\text{Grid}(\mathcal{M})$  and each of its subclasses have rational generating functions.
- ▶ The basis and the generating function for  $\text{Grid}(\mathcal{M})$  can be effectively computed from  $\mathcal{M}$ .
- ▶ The sets of  $\oplus$ -indecomposable,  $\ominus$ -indecomposable and simple permutations in  $\text{Grid}(M)$  all have rational generating functions.



# Out of the woods?

## Conjecture

Every grid class is finitely based.

## Conjecture

Every grid class has an algebraic generating function.

## Question

Are the basis and the generating function for a grid class algorithmically computable from the gridding matrix?

## Question

Is there an algorithm which decides whether a finitely based pattern class (given by its basis) is a grid class?

# Evidence

- ▶ Waton (2007): Grid  $\begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$  is finitely based.
- ▶ Stankova (1994), Atkinson (1998): Grid  $\begin{pmatrix} -1 & 1 \\ 1 & -1 \end{pmatrix}$  has basis  $\{2143, 3412\}$  and generating function

$$\frac{1 - 3x}{(1 - 2x)\sqrt{1 - 4x}}.$$

## Further directions

Generalised grid classes: Allow cells in the matrix to contain arbitrary pattern classes.

V. Vatter, Small permutation classes, submitted.

R. Brignall, Grid classes and partial well order, submitted.



Thank you!

