

Mathematics 101
Fall 2014
Homework 6

1. Let T be a linear operator on a finite dimensional vector space V over a field k , and let $q_1 \mid q_2 \mid \cdots \mid q_s$ be the invariant factors associated to V as a torsion $k[x]$ -module. Show that $q_1 q_2 \cdots q_s = \chi_T$, where χ_T is the characteristic polynomial of T . Show that the minimal polynomial of T divides the characteristic polynomial. Note that this proves the Cayley-Hamilton theorem. *Hint:* You may use without proof (though you should think about it) that the determinant of a matrix of the form $\begin{pmatrix} A & B \\ 0 & C \end{pmatrix}$ with A and C square matrices is the product $\det(A) \det(C)$.
2. Find all rational and Jordan canonical forms of a matrix in $M_5(\mathbb{C})$ having minimal polynomial $x^2(x-1)$. Be sure to give the corresponding invariants and the characteristic polynomials.
3. Show that any linear operator T on a finite dimensional vector space (over a field of characteristic not equal to 2) which satisfies $T^2 = I$ is diagonalizable. Give all possible Jordan forms for 4×4 matrices A with $A^2 = I$.

4. Consider a matrix of the form $A = \begin{pmatrix} \lambda & \mu & 0 & \cdots & 0 \\ 0 & \lambda & \mu & \cdots & 0 \\ 0 & 0 & \ddots & \ddots & 0 \\ 0 & 0 & \cdots & \lambda & \mu \\ 0 & 0 & \cdots & 0 & \lambda \end{pmatrix}$, i.e., with diagonal λ and

μ on the superdiagonal. Find the Jordan canonical form(s) of A . The answer should depend slightly on μ . What does your answer say about the form of Jordan blocks as introduced in the text in comparison with the way we defined them?

5. #24, p501. Prove that there are no 3×3 A matrices over \mathbb{Q} which satisfy $A^8 = I$, but $A^4 \neq I$.
6. #19, p501. Prove that all $n \times n$ matrices over a field F having a fixed characteristic polynomial $f \in F[x]$ are similar if and only if f factors into distinct irreducibles in $F[x]$.