

Mathematics 111
Spring 2009
Homework 2

1. Let R be a ring with identity. Show that the sequence of left R -modules

$$0 \longrightarrow L \xrightarrow{\varphi} M \xrightarrow{\psi} N$$

is exact if and only if for all left R -modules D , the sequence

$$0 \longrightarrow \text{Hom}_R(D, L) \xrightarrow{\varphi_*} \text{Hom}_R(D, M) \xrightarrow{\psi_*} \text{Hom}_R(D, N)$$

is exact.

Hint: We have done the forward direction in class; for the converse, a single propitious choice of D can work, but you still need to sweat the details.

2. Let R be a ring with identity. Show that the sequence of left R -modules

$$L \xrightarrow{\varphi} M \xrightarrow{\psi} N \longrightarrow 0$$

is exact if and only if for all left R -modules D , the sequence

$$0 \longrightarrow \text{Hom}_R(N, D) \xrightarrow{\psi^*} \text{Hom}_R(M, D) \xrightarrow{\varphi^*} \text{Hom}_R(L, D)$$

is exact.

Hint: We have done the forward direction in class. The converse is more complicated than the covariant version; you may want to choose different modules D to establish the various conditions determining exactness of the original sequence. For example, to show ψ is surjective, let $D = N/\text{Im}(\psi)$ (the cokernel of ψ), and $\pi : N \rightarrow D$ the natural projection. Now consider $\psi^*(\pi)$ and its implications.

As a second hint, to show $\text{Im}(\varphi) \subseteq \text{Ker}(\psi)$, you need only show that $\psi \circ \varphi = 0$. Choose $D = N$ and consider the identity map $id_N \in \text{Hom}_R(N, D) = \text{Hom}_R(N, N)$.

3. Let R be a ring with identity and consider the short exact sequence of R -modules:

$$0 \longrightarrow L \xrightarrow{\varphi} M \xrightarrow{\psi} N \longrightarrow 0$$

- Show that if L and N are finitely generated, so is M .
 - Show that if M is finitely generated, so is N .
 - Show by example that if M is finitely generated, L need not be.
4. Determine the number of group homomorphisms $\mathbb{Z}_{12} \oplus \mathbb{Z}_{14} \rightarrow \mathbb{Z}_{20}$, and explicitly characterize them by specifying their action on $(\bar{1}, \bar{0})$ and $(\bar{0}, \bar{1})$.