

# MAJOR FACTS ABOUT DIRECTIONAL DERIVATIVES

- FACT 1. Let  $f : X \subset \mathbb{R}^n \rightarrow \mathbb{R}$  be a function that is differentiable at  $a \in X$ . **Then** the directional derivative  $D_{\mathbf{u}}f(\mathbf{a})$  exists for all unit vectors  $\mathbf{u} \in \mathbb{R}^n$  (that is, for all directions) and  $D_{\mathbf{u}}f(\mathbf{a}) = \nabla f(\mathbf{a}) \cdot \mathbf{u}$ .
- FACT 2. For a fixed  $\mathbf{a} \in \mathbb{R}^n$ , the directional derivative  $D_{\mathbf{u}}f(\mathbf{a})$  is maximized when  $\mathbf{u}$  points in the **same** direction as  $\nabla f(\mathbf{a})$  and is minimized when  $\mathbf{u}$  points in the **opposite** direction. The maximal and minimal values of  $D_{\mathbf{u}}f(\mathbf{a})$  are  $\|\nabla f(\mathbf{a})\|$  and  $-\|\nabla f(\mathbf{a})\|$ , respectively.
- FACT 3. Let  $f : X \subset \mathbb{R}^n \rightarrow \mathbb{R}$  be a differentiable function that has all its partial derivatives continuous,  $S = \{\mathbf{x} \in X \mid f(\mathbf{x}) = c\}$  be a level set for  $f$ , and  $\mathbf{a} \in S$ . **Then**  $\nabla f(\mathbf{a})$  is perpendicular to  $S$ .
- FACT 4. Let  $S \subset \mathbb{R}^3$  be a surface defined by an equation  $f(x, y, z) = c$  and  $\mathbf{x}_0 \in S$ . Assume that  $f$  is differentiable at  $\mathbf{x}_0$  and has all its partial derivatives continuous. **Then** the tangent plane to  $S$  at  $\mathbf{x}_0$  is given by the equation  $\nabla f(\mathbf{x}_0) \cdot (\mathbf{x} - \mathbf{x}_0) = 0$ .