


§7.3 Trig Sub

Motivating example: find area of circle $x^2 + y^2 = 9$



$$\text{Area} = 2 \cdot \int_{-3}^3 \sqrt{9-x^2} dx$$

Problem: we have no method for integrating $y = \sqrt{9-x^2}$
(u-sub does NOT work).

$$x = 3 \sin \theta \quad dx = 3 \cos \theta d\theta$$

$$\sqrt{9-x^2} = \sqrt{9-9\sin^2\theta} = \sqrt{9(1-\sin^2\theta)} = \sqrt{9\cos^2\theta} = 3\cos\theta$$


$$\text{Area} = 2 \cdot \int_{-3}^3 \sqrt{9-x^2} dx = 2 \int_{x=-3}^{x=3} 3\cos\theta \cdot 3\cos\theta d\theta = 18 \int_{x=-3}^{x=3} \cos^2\theta d\theta$$

$$= 18 \cdot \frac{1}{2} (\theta + \sin\theta \cdot \cos\theta) \Big|_{x=-3}^{x=3} = 9 \cdot \left(\theta + \sin\left(\frac{x}{3}\right) + \frac{x}{3} \cdot \frac{\sqrt{9-x^2}}{3} \right) \Big|_{-3}^3$$

$$= 9 \cdot \arcsin(1) - 9 \cdot \arcsin(-1)$$

$$= 9 \cdot \frac{\pi}{2} - 9 \cdot \left(-\frac{\pi}{2}\right)$$

$$= 9\pi$$



$$\sin\theta = \frac{x}{3}$$

$$\text{hence } \theta = \arcsin\left(\frac{x}{3}\right)$$

$$\text{and } \cos\theta = \frac{\sqrt{9-x^2}}{3}$$

Sub calc for

$$\text{ex 1} \int \frac{x^3}{\sqrt{x^2-1}} dx$$

$$x = \sec \theta \quad dx = \sec \theta \tan \theta d\theta$$


$$\sqrt{x^2-1} = \sqrt{\sec^2 \theta - 1} = \sqrt{\tan^2 \theta} = \tan \theta$$

$$= \int \frac{\sec^3 \theta}{\tan \theta} \cdot \sec \theta \cdot \tan \theta d\theta = \int \sec^4 \theta d\theta = \int \sec^2 \theta \cdot \sec^2 \theta d\theta$$

$$= \int (1 + \tan^2 \theta) \cdot \sec^2 \theta d\theta \quad u = \tan \theta \quad du = \sec^2 \theta d\theta$$

$$= \int 1 + u^2 du = u + \frac{u^3}{3} = \tan \theta + \frac{1}{3} \tan^3 \theta$$

$$= \sqrt{x^2-1} + \frac{1}{3} (x^2-1)^{3/2} + C$$



$$\text{ex 1} \int \frac{x-1}{\sqrt{9-x^2}} dx$$

$$x = 3 \sin \theta \quad dx = 3 \cos \theta d\theta$$

$$= \int \frac{3 \sin \theta - 1}{3 \cos \theta} \cdot 3 \cos \theta d\theta = \int 3 \sin \theta - 1 d\theta = -3 \cos \theta - \theta$$

$$= -3 \cdot \frac{\sqrt{9-x^2}}{3} - \arcsin\left(\frac{x}{3}\right) + C$$

$$= -\sqrt{9-x^2} - \arcsin\left(\frac{x}{3}\right) + C$$




Table of Trig Sub

expression	substitution	Pythagorus
$\sqrt{a^2 - x^2}$	$x = a \sin \theta$	$a^2 - a^2 \sin^2 \theta = a^2 \cos^2 \theta$
$\sqrt{a^2 + x^2}$	$x = a \tan \theta$	$a^2 + a^2 \tan^2 \theta = a^2 \sec^2 \theta$
$\sqrt{x^2 - a^2}$	$x = a \sec \theta$	$a^2 \sec^2 \theta - a^2 = a^2 \tan^2 \theta$