

## The Fundamental Theorem of Calculus

©2002, 2008 Donald Kreider and Dwight Lahr

We are about to discuss a theorem that relates derivatives and definite integrals. It is so important in the study of calculus that it is called the *Fundamental Theorem of Calculus*. It also gives us a practical way to evaluate many definite integrals without resorting to the limit definition. The theorem has two main parts that we will state separately as Part I and Part II.

**Fundamental Theorem of Calculus (Part I-antiderivative):** Suppose that  $f$  is a continuous function on the interval  $I$  containing the point  $a$ . Define the function  $F$  on  $I$  by the integral formula

$$F(x) = \int_a^x f(t) dt$$

Then  $F$  is differentiable on  $I$  and  $F'(x) = f(x)$ . That is,  $F$  is an antiderivative of  $f$  on  $I$ .

**Fundamental Theorem of Calculus (Part II-evaluation):** If  $G(x)$  is any antiderivative of  $f$  on  $I$  (that is,  $G'(x) = f(x)$  on  $I$ ), then for any  $b$  in  $I$ ,

$$\int_a^b f(x) dx = G(b) - G(a)$$

This theorem is truly remarkable. Leibniz seems to have been the first one to recognize its generality and significance. Let's look at some examples so that we can gain a better understanding of what the theorem says, and then we will outline a proof.

**Example 1:** To compute  $\int_0^1 (x+1) dx$ , we need only find an antiderivative of  $x+1$ , namely,  $x^2/2+x$ . Then we evaluate this antiderivative at 1 and subtract its value at 0. Thus,  $\int_0^1 (x+1) dx = (1/2+1) - (0) = 3/2$ .

We normally use a vertical bar to indicate evaluation of the antiderivative at the endpoints of the interval. That is,

$$G(x)|_a^b = G(b) - G(a)$$

**Example 2:**  $\int_0^1 x^2 dx = \left. \frac{x^3}{3} \right|_0^1 = \frac{1}{3} - 0 = \frac{1}{3}$ .

**Example 3:**  $\int_0^{\pi/4} \sin x dx = -\cos x|_0^{\pi/4} = -1/\sqrt{2} - (-1) = 1 - 1/\sqrt{2}$ .

**Example 4:**  $\int_0^{\pi/4} \sec^2 x dx = \tan x|_0^{\pi/4} = 1 - 0 = 1$ .

We can also illustrate Part I of the Fundamental Theorem.

**Example 5:**  $\frac{d}{dx} \int_1^x t^2 dt = x^2$ .

**Example 6:**  $\frac{d}{dx} \int_1^{x^2} t^3 dt = (x^2)^3 \cdot 2x$  where we first have used the Fundamental Theorem and then the chain rule to complete the calculation of the derivative.

**Example 7:** Consider  $\frac{d}{dx} \int_{x^2}^{x^3} e^{-t^2} dt$ . We first have to put the integral in the correct form so that we can use the Fundamental Theorem:


$$\begin{aligned} \frac{d}{dx} \int_{x^2}^{x^3} e^{-t^2} dt &= \\ &= \frac{d}{dx} \left( \int_{x^2}^0 e^{-t^2} dt + \int_0^{x^3} e^{-t^2} dt \right) \\ &= \frac{d}{dx} \left( -\int_0^{x^2} e^{-t^2} dt + \int_0^{x^3} e^{-t^2} dt \right) \\ &= -e^{-x^4} (2x) + e^{-x^6} (3x^2) \end{aligned}$$

Now that we have gained some experience with the Fundamental Theorem through examples, let's look at a sketch of a proof in a special case.

**Proof of the Fundamental Theorem (Part I):** Fix  $x$  in  $I$ . Given that  $F(x) = \int_a^x f(t) dt$ , we need to evaluate the limit

$$\lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h}$$

But look at the sketch below. Notice that  $F(x)$  is the area under the graph of  $f$  and above the interval  $[a, x]$ , while  $F(x+h)$  is the area under the graph of  $f$  and above the interval  $[a, x+h]$ . Thus,  $F(x+h) - F(x)$  is the area under the graph of  $f$  and above the interval  $[x, x+h]$ .



But for small values of  $h$ , this area is approximately equal to the area of the rectangle of height  $f(x)$  on the same base; its area is length times width, or  $h \cdot f(x)$ . Thus, for small  $h$ , the difference quotient is approximately equal to  $\frac{hf(x)}{h} = f(x)$ . In other words,

$$F'(x) = \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} = f(x)$$

thereby completing the proof of Part I.

**Proof of the Fundamental Theorem (Part II):** From Part I, we have that  $F(x) = \int_a^x f(t) dt$  is an antiderivative of  $f$ . If  $G$  is another antiderivative, then we know from a previous result that they must differ by a constant. That is,  $G(x) = F(x) + C$ . Now, we know that  $F(a) = \int_a^a f(t) dt = 0$ . Thus, we can determine the value of  $C$ :  $G(a) = F(a) + C = 0 + C = C$ . Hence,  $G(x) = F(x) + G(a)$ , or  $F(x) = G(x) - G(a)$ . So, if  $b$  is any point in  $I$ , we have  $G(b) - G(a) = F(b) = \int_a^b f(t) dt$ , which is what we wanted to prove.

### Another Proof of the Fundamental Theorem of Calculus

**Theorem statement:** If  $G(x)$  is any antiderivative of  $f$  on  $I$  (that is,  $G'(x) = f(x)$  on  $I$ ), then for any  $b$  in  $I$ ,

$$\int_a^b f(x) dx = G(b) - G(a)$$

We are going to prove this result by an application of Euler's Method which we studied earlier. Suppose we consider the Initial Value Problem

$$\text{IVP: } y' = f(x), y(a) = 0, a \leq x \leq b, \text{ where } a, b \text{ are in } I,$$

and we want to find  $y(b)$ . Then because both  $y$  and  $G$  are antiderivatives of  $f$  on  $[a, b]$ ,  $y(x) = G(x) + C$  for some constant  $C$  on  $[a, b]$ . Then  $0 = y(a) = G(a) + C$  implies  $C = -G(a)$  and hence  $y(b) = G(b) - G(a)$ . Now, we will use Euler's method to approximate  $y(b)$ .

Suppose we use an integral number  $n$  of steps where each step has size  $\frac{b-a}{n}$ . Then, starting at the point  $(a, 0)$  where the slope is  $y'(a) = f(a)$ , we generate the following points:

Point $(x, y)$	Slope
$(a, 0)$	$f(a)$
$(a + h, f(a) h)$	$f(a + h)$
$(a + 2h, f(a) h + f(a + h) h)$	$f(a + 2h)$
etc.	etc.

The endpoint at  $x = b$  has  $y$ -coordinate

$$\sum_{i=0}^{n-1} f(a + ih) h$$

The above sum is the Euler method approximate value of  $y(b)$  which converges to  $y(b)$  as  $h \rightarrow 0$ . But note that it is also a Riemann sum for the definite integral from  $a$  to  $b$  of  $f$ , and the Riemann sum converges to the value of the integral as  $h \rightarrow 0$ . Thus, because the limit of the sum is unique, we have

$$y(b) = \int_a^b f(x) dx$$

and from the result  $y(b) = G(b) - G(a)$  in the first paragraph of the proof, we see that the proof is complete.

**Exercises:** [Problems](#) **Check what you have learned!**

**Videos:** [Tutorial Solutions](#) **See problems worked out!**