
Limits

From last time...

Let y = f(t) be a function that gives the position at time t of an object moving along the y-axis. Then

Ave
$$\operatorname{vel}[t_1, t_2] = \frac{f(t_2) - f(t_1)}{t_2 - t_1}$$

$$\operatorname{Velocity}(t) = \lim_{h \to 0} \frac{f(t+h) - f(t)}{h}.$$




We need to be able to take limits!

Limit of a Function – Definition

We say that a function f approaches the limit L as x approaches a,


written
$$\lim_{x \to a} f(x) = L$$
,

if we can make f(x) as close to L as we want by taking x sufficiently close to a.



i.e. If you need Δy to be smaller, you only need to make Δx smaller (Δ means "change")

One-sided limits



Right-handed limit: $L_r = \lim_{x \to a^+} f(x)$

if f(x) gets closer to L_r as x gets closer to a from the right

Left-handed limit: $L_{\ell} = \lim_{x \to a^{-}} f(x)$

if f(x) gets closer to L_{ℓ} as x gets closer to a from the left

Theorem

The limit of f as $x \to a$ exists if and only if both the right-hand and left-hand limits exist and have the same value, i.e.

$$\lim_{x\to a} f(x) = L \text{ if and only if } \lim_{x\to a^-} f(x) = L \text{ and } \lim_{x\to a^+} f(x) = L.$$

Examples

$$\lim_{x \to 2} \frac{x-2}{x+3} \qquad \lim_{x \to 1} \frac{x^2-1}{x-1} \qquad \lim_{x \to 0} \frac{1}{x}$$

Theorem

If $\lim_{x\to a} f(x) = A$ and $\lim_{x\to a} g(x) = B$ both exist, then

- 1. $\lim_{x\to a} (f(x) + g(x)) = \lim_{x\to a} f(x) + \lim_{x\to a} g(x) = A + B$
- 2. $\lim_{x\to a} (f(x) g(x)) = \lim_{x\to a} f(x) \lim_{x\to a} g(x) = A B$
- 3. $\lim_{x\to a} (f(x)g(x)) = \lim_{x\to a} f(x) \cdot \lim_{x\to a} g(x) = A \cdot B$
- 4. $\lim_{x\to a} (f(x)/g(x)) = \lim_{x\to a} f(x)/\lim_{x\to a} g(x) = A/B \ (B \neq 0).$

In short: to take a limit

- Step 1: Can you just plug in? If so, do it.
- Step 2: If not, is there some sort of algebraic manipulation (like cancellation) that can be done to fix the problem? If so, do it. Then plug in.
- Step 3: Learn some special limit to fix common problems. (Later) If in doubt, graph it!

Examples

1.
$$\lim_{x\to 2} \frac{x-2}{x+3} = \boxed{0}$$
 because if $f(x) = \frac{x-2}{x+3}$, then $f(2) = 0$.

2.
$$\lim_{x \to 1} \frac{x^2 - 1}{x - 1} \xrightarrow[\to 0]{}_{\to 0}$$

If $f(x) = \frac{x^2-1}{x-1}$, then f(x) is undefined at x = 1. However, so long as $x \neq 1$,

$$f(x) = \frac{x^2 - 1}{x - 1} = \frac{(x + 1)(x - 1)}{x - 1} = x + 1.$$

So

$$\lim_{x \to -1} \frac{x^2 - 1}{x - 1} = \lim_{x \to 1} x + 1 = 1 + 1 = 2.$$

3. $\lim_{x\to 0} \frac{\sqrt{x+2}-\sqrt{2}}{x} \xrightarrow{\to 0}$, so again, f(x) is undefined at a.

Examples

3. $\lim_{x\to 0} \frac{\sqrt{x+2}-\sqrt{2}}{x} \xrightarrow[]{\to 0}$, so again, f(x) is undefined at a.

Multiply top and bottom by the conjugate:

$$\lim_{x \to 0} \frac{\sqrt{x+2} - \sqrt{2}}{x} = \lim_{x \to 0} \left(\frac{\sqrt{x+2} - \sqrt{2}}{x} \right) \left(\frac{\sqrt{x+2} + \sqrt{2}}{\sqrt{x+2} + \sqrt{2}} \right)$$

$$= \lim_{x \to 0} \frac{x+2-2}{x(\sqrt{x+2} + \sqrt{2})} \quad \text{since } (a-b)(a+b) = a^2 - b^2$$

$$= \lim_{x \to 0} \frac{x}{x(\sqrt{x+2} + \sqrt{2})}$$

$$= \lim_{x \to 0} \frac{1}{\sqrt{x+2} + \sqrt{2}} = \boxed{\frac{1}{2\sqrt{2}}}$$

Examples

1.
$$\lim_{x \to 1} \frac{x^2 - 3x + 2}{x^2 + 4x - 5}$$

$$2. \lim_{x \to -2} \frac{|x|}{x}$$

3.
$$\lim_{x\to 0} \frac{|x|}{x}$$


4.
$$\lim_{h \to 0} \frac{(3+x)^2 - 3^2}{x}$$

Infinite limits

If f(x) gets arbitrarily large as $x \to a$, then it doesn't have a limit. Sometimes, though, it's more useful to give more information.

Example:

For both $f(x) = \frac{1}{x}$ and $f(x) = \frac{1}{x^2}$, $\lim_{x\to 0} f(x)$ does not exist. However, they're both "better behaved" than that might imply:




Why? A vertical asymptote occurs where

$$\lim_{x \to a^+} f(x) = \pm \infty$$
 and $\lim_{x \to a^-} f(x) = \pm \infty$

Infinite limits

Badly behaved example:





(denser and denser vertical asymptotes)

 $\lim_{x \to a^+} \csc(1/x)$ does not exist, and $\lim_{x \to a^-} \csc(1/x)$ does not exist


Limits at Infinity

We say that a function f approaches the limit L as x gets bigger and bigger (in the positive or negative direction),

written
$$\lim_{x \to \infty} f(x) = L$$
 or $\lim_{x \to -\infty} f(x) = L$

if we can make f(x) as close to L as we want by taking x sufficiently large. (By "large", we mean large in magnitude)

Example 1:




$$\lim_{x \to -\infty} \frac{1}{x} = 0 \qquad \text{and} \qquad \lim_{x \to \infty} \frac{1}{x} = 0.$$

Limits at Infinity: functions and their inverses

Theorem

If
$$\lim_{x\to a^{\pm}} f(x) = \infty$$
, then $\lim_{x\to\infty} f^{-1}(x) = a$.
If $\lim_{x\to a^{\pm}} f(x) = -\infty$, then $\lim_{x\to -\infty} f^{-1}(x) = a$.

Example: Let arctan(x) be the inverse function to tan(x):



Since
$$\lim_{x \to \pi/2^-} = \infty$$
 and $\lim_{x \to -\pi/2^+} = -\infty$

(restrict the domain to $\left(-\frac{\pi}{2},\frac{\pi}{2}\right)$ to that we can invert)

$$y = \arctan(x):$$

$$\xrightarrow{\pi/2}$$

$$\xrightarrow{-\pi/2}$$

we have
$$\lim_{x\to\infty} = \pi/2$$
 and $\lim_{x\to-\infty} = -\pi/2$

Rational functions

Limits that look like they're going to $\frac{\infty}{\infty}$ can actually be doing lots of different things. To fix this, divide the top and bottom by the highest power in the denominator!

Ex 1
$$\lim_{x \to \infty} \frac{x-1}{x^3+2} \xrightarrow{\to \infty}$$

Fix: multiply the expression by $\frac{1/x^3}{1/x^3}$:

$$\lim_{x \to \infty} \frac{x-1}{x^3+2} = \lim_{x \to \infty} \left(\frac{x-1}{x^3+2}\right) \left(\frac{1/x^3}{1/x^3}\right) = \lim_{x \to \infty} \frac{\frac{1}{x^2} - \frac{1}{x^3}}{1 + \frac{2}{x^3}} = \frac{0-0}{1+0} = \boxed{0}$$

Ex 2
$$\lim_{x\to\infty} \frac{3x^2 - 2x + 1}{4x^2 - 1} \xrightarrow[]{}^{+\infty}$$

Fix: multiply the expression by $\frac{1/x^2}{1/x^2}$

Ex 3
$$\lim_{x \to \infty} \frac{x^4 - x^2 + 2}{x^3 + 3} \xrightarrow{\infty} \infty$$


Fix: multiply the expression by $\frac{1/x^3}{1/x^3}$


Rational functions: quick trick


$$\lim_{x \to \infty} \frac{x - 1}{x^3 + 2} = 0$$

$$\lim_{x \to \infty} \frac{3x^2 - 2x + 1}{4x^2 - 1} = \frac{3}{4}$$

$$\lim_{x \to \infty} \frac{x - 1}{x^3 + 2} = 0 \qquad \lim_{x \to \infty} \frac{3x^2 - 2x + 1}{4x^2 - 1} = \frac{3}{4} \qquad \lim_{x \to \infty} \frac{x^4 - x^2 + 2}{x^3 + 3} = \infty$$







Suppose

$$P(x) = a_n x^n + \dots + a_1 x + a_0$$
 and $Q(x) = b_m x^m + \dots + b_1 x + b_0$

$$Q(x) = b_m x^m + \cdots + b_1 x + b_0$$

are polynomials of degree n and m respectively. Then in general,

$$\lim_{x \to \infty} \frac{P(x)}{Q(x)} = \lim_{x \to \infty} \frac{a_n x^n}{b_m x^m} = \lim_{x \to \infty} \frac{a_n}{b_m} x^{n-m} = \begin{cases} 0 & n < m \\ \frac{a_n}{b_m} & n = m \\ \pm \infty & n < m \end{cases}$$

Examples: Other ratios with powers.

Example:

$$\lim_{x \to \infty} \frac{x}{\sqrt{3x^2 + 2}}$$

[hint: multiply by $\frac{1/x}{1/x}$ and remember $a\sqrt{b}=\sqrt{a^2b}$.]

Evaluating limits when $x \to 0$.

1. Show
$$\lim_{x\to 0} (x^2 - 2)^2 + 6 = 10$$
.

2. Show
$$\lim_{x \to 0} \frac{5x}{x} = 5$$
.

3. Show
$$\lim_{x\to 0} \frac{17x}{2x} = 17/2$$
.

4. Show
$$\lim_{x\to 0} \frac{-317x}{422x} = -317/422$$
.

5. Show
$$\lim_{x\to 0} \frac{-317x - 3}{422x + 5} = -3/5$$
.

6. Show
$$\lim_{h\to 0} \frac{\sqrt{x+h} - \sqrt{x}}{h} = \frac{1}{2\sqrt{x}}.$$

7. Show
$$\lim_{x\to 0} \frac{\sqrt{1+x+x^2}-1}{x} = 1/2$$
.

8. Show
$$\lim_{x\to 0} \frac{\sqrt{3+x} - \sqrt{3}}{x} = 1/(2\sqrt{3}).$$

9. Show
$$\lim_{h\to 0} \frac{1}{h} \left(\frac{1}{\sqrt{x+h}} - \frac{1}{\sqrt{x}} \right) = -(1/2)x^{-3/2}$$
.

10. Show
$$\lim_{x\to 0} \frac{2x}{\sqrt{a+x} - \sqrt{a-x}} = 2\sqrt{a}$$
.

11. Show
$$\lim_{x\to 0} \frac{\sqrt{1+x}-1}{x} = 1/2$$
.

12. Show
$$\lim_{x\to 0} \frac{x}{\sqrt{1+x}-1} = 2$$
.

13. Show
$$\lim_{x\to 0} \frac{e^x + e^{-x} - 2}{x^2} = 1$$
.

14. Show

$$\lim_{h \to 0} \frac{f(x+h) - f(x)}{h} = \frac{a}{2\sqrt{ax+b}}$$

when
$$f(x) = \sqrt{ax + b}$$
.

15. Show

$$\lim_{h \to 0} \frac{f(x+h) - f(x)}{h} = mn(mx+c)^{n-1}$$

when $f(x) = (mx + c)^n$.

Evaluating limits when $x \to a$.

1. Show
$$\lim_{x \to 1} (6x^2 - 4x + 3) = 5$$
.

2. Show
$$\lim_{x \to 7} \frac{x^2 - 49}{x - 7} = 14$$
.

3. Show
$$\lim_{x \to 2} \frac{x^2 - 6x + 8}{x - 2} = -2$$
.

4. Show
$$\lim_{x \to -5} \frac{2x^2 + 9x - 5}{x + 5} = -11$$
.

5. Show
$$\lim_{x \to 1} \frac{x^3 - 1}{x - 1} = 3$$
.

6. Show
$$\lim_{x\to 3} \frac{x^2 - 4x + 3}{x^2 - 2x - 3} = 1/2$$
.

7. Show
$$\lim_{x \to -2} \frac{x^3 + 8}{x + 2} = 4$$
.

8. Show
$$\lim_{x \to 3} \frac{x^4 - 81}{x - 3} = 108$$
.

9. Show
$$\lim_{x \to 5} \frac{x^5 - 3125}{x - 5} = 3125$$
.

10. Show
$$\lim_{x \to a} \frac{x^{12} - a^{12}}{x - a} = 12a^{11}$$
.

11. Show
$$\lim_{x \to a} \frac{x^{5/2} - a^{5/2}}{x - a} = (5/2)a^{3/2}$$
.

12. Show
$$\lim_{x \to a} \frac{(x+2)^{5/3} - (a+2)^{5/3}}{x-a} = (5/3)(a+2)^{2/3}$$
.

13. Show
$$\lim_{x \to 4} \frac{x^3 - 64}{x^2 - 16} = 6$$
.

14. Show
$$\lim_{x \to 2} \frac{x^5 - 32}{x^3 - 8} = 20/3$$
.

15. Show
$$\lim_{x \to 1} \frac{x^n - 1}{x - 1} = n$$
.

16. Show
$$\lim_{x \to a} \frac{\sqrt{x} - \sqrt{a}}{x - a} = \frac{1}{2\sqrt{a}}.$$

17. Show
$$\lim_{x\to 2} \frac{\sqrt{3-x}-1}{2-x} = 1/2$$
.

18. Show
$$\lim_{x \to a} \frac{\sqrt{a+2x} - \sqrt{3x}}{\sqrt{3a+x} - 2\sqrt{x}} = \frac{2\sqrt{3}}{9}$$
.

19. Show
$$\lim_{x \to a} \frac{x^n - a^n}{x - a} = na^{n-1}$$
.

Evaluating limits when $x \to \infty$.

1. Show
$$\lim_{x \to \infty} \frac{x+2}{x-2} = 1$$
.

6. Show
$$\lim_{x \to \infty} \frac{x}{\sqrt{4x^2 + 1} - 1} = 1/2$$
.

2. Show
$$\lim_{x\to\infty} \frac{3x^2 + 2x - 5}{5x^2 + 3x + 1} = 3/5$$
.

7. Show
$$\lim_{x \to -\infty} 2^x = 0$$
.

3. Show
$$\lim_{x\to\infty} \frac{x^2 - 7x + 11}{3x^2 + 10} = 1/3$$
.

8. Show
$$\lim_{t \to \infty} \frac{t+1}{t^2+1} = 0$$
.

4. Show
$$\lim_{x \to \infty} \frac{2x^3 - 5x + 7}{7x^3 + 2x^2 - 6} = 2/7$$
.

9. Show
$$\lim_{n \to \infty} \sqrt{n^2 + 1} - n = 0$$
.

5. Show
$$\lim_{x \to \infty} \frac{(3x-1)(4x-5)}{(x+6)(x-3)} = 12.$$

10. Show
$$\lim_{n \to \infty} \sqrt{n^2 + n} - n = 1/2$$
.