


Continuity

Warm-up

Suppose the graph of $y = f(x)$ looks like



answers to 4.:


a	$x \rightarrow a^-$	$x \rightarrow a^+$

1. What is the domain of $f(x)$?
2. What is the range of $f(x)$?
3. For which values a in $[1, 7]$ does $\lim_{x \rightarrow a} f(x)$ not exist?
4. For those values you picked out in 3., what are $\lim_{x \rightarrow a^-} f(x)$ and $\lim_{x \rightarrow a^+} f(x)$?
5. Which values a satisfy

$$f(a) \text{ and } \lim_{x \rightarrow a} f(x) \text{ exist, but } f(a) \neq \lim_{x \rightarrow a} f(x)?$$

Warm-up

Suppose the graph of $y = f(x)$ looks like



answers to 4.:

a	$x \rightarrow a^-$	$x \rightarrow a^+$
1	DNE	-1
4	2	1
6	1.5	2
7	3	DNE

1. What is the domain of $f(x)$?
2. What is the range of $f(x)$?
3. For which values a in $[1, 7]$ does $\lim_{x \rightarrow a} f(x)$ not exist?

$$(1, 2) \cup (2, 6) \cup (6, 7]$$

$$[-1, 3]$$

4. For those values you picked out in 3., what are

$$\lim_{x \rightarrow a^-} f(x) \quad \text{and} \quad \lim_{x \rightarrow a^+} f(x)?$$

$$a = 1, 4, 6, 7$$


(see above)

5. Which values a satisfy

$$a = 5$$

$$f(a) \text{ and } \lim_{x \rightarrow a} f(x) \text{ exist, but } f(a) \neq \lim_{x \rightarrow a} f(x)?$$


Domain definitions



Let D be the domain of $f(x)$.

$$\text{Ex. } D = (1, 2) \cup (2, 6) \cup (6, 7]$$

Domain definitions



Let D be the domain of $f(x)$.


$$\text{Ex. } D = (1, 2) \cup (2, 6) \cup (6, 7]$$

Definition

An *interior point* of D is any point **in** D which is not an endpoint or an isolated point.

Ex. Everything in D except $x = 7$.

Domain definitions



Let D be the domain of $f(x)$.

Ex. $D = (1, 2) \cup (2, 6) \cup (6, 7]$

Ex 2. $D = \{\frac{1}{2}\} \cup (1, 2) \cup (2, 6) \cup (6, 7]$

Definition


An *interior point* of D is any point **in** D which is not an endpoint or an isolated point.

Ex. Everything in D except $x = 7$.

Ex 2. Everything in D except $x = \frac{1}{2}$ & 7.

Continuity

Let a be an interior point of D .




Definition

A function is *continuous* at a if $\lim_{x \rightarrow a} f(x) = f(a)$. If it is not continuous at a , then function is *discontinuous* at a .

Continuity

Let a be an interior point of D .



Definition

A function is *continuous* at a if $\lim_{x \rightarrow a} f(x) = f(a)$. If it is not continuous at a , then function is *discontinuous* at a .


Checklist:

1. Is a an interior point? *If no, stop here... we'll get back to these.*
2. Does (a) $\lim_{x \rightarrow a^-} f(x)$ exist? (b) $\lim_{x \rightarrow a^+} f(x)$ exist?
3. Does $\lim_{x \rightarrow a} f(x)$ exist? (i.e. does (a) = (b)?)
4. Does $f(a) = \lim_{x \rightarrow a} f(x)$?

If the answer to any of 2.-4. is "no", then $f(x)$ is discontinuous at a .

Continuity

Let a be an interior point of D .



Ex. $f(x)$ is discontinuous
at $x = 4$ and 5 .
No other points are fair game!

Definition

A function is *continuous* at a if $\lim_{x \rightarrow a} f(x) = f(a)$. If it is not continuous at a , then function is *discontinuous* at a .

Checklist:

1. Is a an interior point? If no, stop here... we'll get back to these.
2. Does (a) $\lim_{x \rightarrow a^-} f(x)$ exist? (b) $\lim_{x \rightarrow a^+} f(x)$ exist?
3. Does $\lim_{x \rightarrow a} f(x)$ exist? (i.e. does (a) = (b)?)
4. Does $f(a) = \lim_{x \rightarrow a} f(x)$?

If the answer to any of 2.-4. is "no", then $f(x)$ is discontinuous at a .

Some examples:

Over their domains, all

polynomials, rational functions, trigonometric functions,
exponential functions, absolute values,

and their inverses are all continuous functions.

(Jumps all happen over domain gaps)

Some examples:

Over their domains, all

polynomials, rational functions, trigonometric functions,
exponential functions, absolute values,

and their inverses are all continuous functions.

(Jumps all happen over domain gaps)

Example: Is the function $f(x) = \begin{cases} x^2 & x < 1 \\ x^3 + 2 & 1 \leq x \end{cases}$ continuous?

Some examples:

Over their domains, all

polynomials, rational functions, trigonometric functions,
exponential functions, absolute values,

and their inverses are all continuous functions.

(Jumps all happen over domain gaps)

Example: Is the function $f(x) = \begin{cases} x^2 & x < 1 \\ x^3 + 2 & 1 \leq x \end{cases}$ continuous?

Solution: The only possible problem would happen at $x = 1$. Let's check there:

Some examples:

Over their domains, all

polynomials, rational functions, trigonometric functions,
exponential functions, absolute values,

and their inverses are all continuous functions.

(Jumps all happen over domain gaps)

Example: Is the function $f(x) = \begin{cases} x^2 & x < 1 \\ x^3 + 2 & 1 \leq x \end{cases}$ continuous?

Solution: The only possible problem would happen at $x = 1$. Let's check there:

$$\lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^-} x^2$$

$$\lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1^+} x^3 + 2$$

Some examples:

Over their domains, all

polynomials, rational functions, trigonometric functions,
exponential functions, absolute values,

and their inverses are all continuous functions.

(Jumps all happen over domain gaps)

Example: Is the function $f(x) = \begin{cases} x^2 & x < 1 \\ x^3 + 2 & 1 \leq x \end{cases}$ continuous?

Solution: The only possible problem would happen at $x = 1$. Let's check there:

$$\lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^-} x^2 = 1$$

$$\lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1^+} x^3 + 2 = 3$$

No, $f(x)$ is discontinuous at $x = 1$ because 1 is an interior point of the domain, but $\lim_{x \rightarrow 1} f(x)$ does not exist.

Some examples:

Over their domains, all

polynomials, rational functions, trigonometric functions,
exponential functions, absolute values,

and their inverses are all continuous functions.


(Jumps all happen over domain gaps)

Example: Is the function $f(x) = \begin{cases} x^2 & x < 1 \\ x^3 + 2 & 1 \leq x \end{cases}$ continuous?

Solution: The only possible problem would happen at $x = 1$. Let's check there:

$$\lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^-} x^2 = 1$$

$$\lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1^+} x^3 + 2 = 3$$



No, $f(x)$ is discontinuous at $x = 1$ because 1 is an interior point of the domain, but $\lim_{x \rightarrow 1} f(x)$ does not exist.

Right Continuity and Left Continuity

Definition

A function $f(x)$ is *right continuous* at a point a if it is defined on an interval $[a, b)$ and $\lim_{x \rightarrow a^+} f(x) = f(a)$.

Similarly, a function $f(x)$ is *left continuous* at a point a if it is defined on an interval $(b, a]$ and $\lim_{x \rightarrow a^-} f(x) = f(a)$.


Right Continuity and Left Continuity

Definition

A function $f(x)$ is *right continuous* at a point a if it is defined on an interval $[a, b)$ and $\lim_{x \rightarrow a^+} f(x) = f(a)$.

Similarly, a function $f(x)$ is *left continuous* at a point a if it is defined on an interval $(b, a]$ and $\lim_{x \rightarrow a^-} f(x) = f(a)$.

Example:




$f(x)$ is

- (a) continuous at every *interior* point in D except $x = 4$ and 5 ;
- (b) only right continuous at those points included in (a); and
- (c) additionally left continuous at $x = 4$ and $x = 7$.

Suppose a function f has no isolated points in its domain.

Definition

A function f is *continuous over its domain D* if **(1)** it is continuous at every interior point of D , and **(2)** it is left (or right) continuous at every endpoint of D . Otherwise, it has a *discontinuity* at each point in D which violates (1) or (2).



Filling and Fixing

Suppose a is a point of discontinuity in D

(a) If a is an interior point and $\lim_{x \rightarrow a} f(x) = L$ exists; or

(b) if a is an endpoint and $\lim_{x \rightarrow a^\pm} f(x) = L$ exists,

then we say $f(x)$ has a *removable discontinuity*:

$$\bar{f}(x) = \begin{cases} f(x) & x \neq a \\ L & x = a \end{cases}$$

Filling and Fixing


Suppose a is a point of discontinuity in D

(a) If a is an interior point and $\lim_{x \rightarrow a} f(x) = L$ exists; or

(b) if a is an endpoint and $\lim_{x \rightarrow a^\pm} f(x) = L$ exists,

then we say $f(x)$ has a *removable discontinuity*:

$$\bar{f}(x) = \begin{cases} f(x) & x \neq a \\ L & x = a \end{cases}$$



Filling and Fixing


Suppose a is a point of discontinuity in D

(a) If a is an interior point and $\lim_{x \rightarrow a} f(x) = L$ exists; or

(b) if a is an endpoint and $\lim_{x \rightarrow a^\pm} f(x) = L$ exists,

then we say $f(x)$ has a *removable discontinuity*:

$$\bar{f}(x) = \begin{cases} f(x) & x \neq a \\ L & x = a \end{cases}$$



Filling and Fixing


Suppose a is a point of discontinuity in D

(a) If a is an interior point and $\lim_{x \rightarrow a} f(x) = L$ exists; or

(b) if a is an endpoint and $\lim_{x \rightarrow a^\pm} f(x) = L$ exists,

then we say $f(x)$ has a *removable discontinuity*:

$$\bar{f}(x) = \begin{cases} f(x) & x \neq a \\ L & x = a \end{cases}$$



Filling and Fixing


Suppose a is a point of discontinuity in D

(a) If a is an interior point and $\lim_{x \rightarrow a} f(x) = L$ exists; or

(b) if a is an endpoint and $\lim_{x \rightarrow a^\pm} f(x) = L$ exists,

then we say $f(x)$ has a *removable discontinuity*:

$$\bar{f}(x) = \begin{cases} f(x) & x \neq a \\ L & x = a \end{cases}$$



Example: $f(x)$ has a removable discontinuity in exactly one place:

$$\bar{f}(x) = \begin{cases} f(x) & x \neq 5 \\ 1 & x = 5 \end{cases}$$

Filling and Fixing

Suppose a is a hole in D (a is arbitrarily close to points in D , but not in D).

(a) If a would be an interior point and $\lim_{x \rightarrow a} f(x) = L$ exists; or

(b) if a would be an endpoint and $\lim_{x \rightarrow a^\pm} f(x) = L$ exists,

then we say $f(x)$ has a *continuous extension*:

$$\bar{f}(x) = \begin{cases} f(x) & x \neq a \\ L & x = a \end{cases}$$

Filling and Fixing


Suppose a is a hole in D (a is arbitrarily close to points in D , but not in D).

(a) If a would be an interior point and $\lim_{x \rightarrow a} f(x) = L$ exists; or

(b) if a would be an endpoint and $\lim_{x \rightarrow a^\pm} f(x) = L$ exists,

then we say $f(x)$ has a *continuous extension*:

$$\bar{f}(x) = \begin{cases} f(x) & x \neq a \\ L & x = a \end{cases}$$



Filling and Fixing


Suppose a is a hole in D (a is arbitrarily close to points in D , but not in D).

(a) If a would be an interior point and $\lim_{x \rightarrow a} f(x) = L$ exists; or

(b) if a would be an endpoint and $\lim_{x \rightarrow a^\pm} f(x) = L$ exists,

then we say $f(x)$ has a *continuous extension*:

$$\bar{f}(x) = \begin{cases} f(x) & x \neq a \\ L & x = a \end{cases}$$



Filling and Fixing


Suppose a is a hole in D (a is arbitrarily close to points in D , but not in D).

(a) If a would be an interior point and $\lim_{x \rightarrow a} f(x) = L$ exists; or

(b) if a would be an endpoint and $\lim_{x \rightarrow a^\pm} f(x) = L$ exists,

then we say $f(x)$ has a *continuous extension*:

$$\bar{f}(x) = \begin{cases} f(x) & x \neq a \\ L & x = a \end{cases}$$



Filling and Fixing


Suppose a is a hole in D (a is arbitrarily close to points in D , but not in D).

(a) If a would be an interior point and $\lim_{x \rightarrow a} f(x) = L$ exists; or

(b) if a would be an endpoint and $\lim_{x \rightarrow a^\pm} f(x) = L$ exists,

then we say $f(x)$ has a *continuous extension*:

$$\bar{f}(x) = \begin{cases} f(x) & x \neq a \\ L & x = a \end{cases}$$



Filling and Fixing


Suppose a is a hole in D (a is arbitrarily close to points in D , but not in D).

(a) If a would be an interior point and $\lim_{x \rightarrow a} f(x) = L$ exists; or

(b) if a would be an endpoint and $\lim_{x \rightarrow a^\pm} f(x) = L$ exists,

then we say $f(x)$ has a *continuous extension*:

$$\bar{f}(x) = \begin{cases} f(x) & x \neq a \\ L & x = a \end{cases}$$



Example: $f(x)$ has continuous extensions in exactly two places:

$$\bar{f}_1(x) = \begin{cases} f(x) & x \neq 1 \\ -1 & x = 1 \end{cases} \quad \text{and} \quad \bar{f}_2(x) = \begin{cases} f(x) & x \neq 2 \\ 1 & x = 2 \end{cases}$$

Examples

- (A) Which of the following have removable discontinuities? For those which do, what are the alternate functions with those discontinuities removed?
- (B) Which of the following have continuous extensions? For those which do, what are those extensions?

1. $f(x) = \frac{x^2 - 4}{x - 2}$

2. $f(x) = \begin{cases} \sin x & x \neq \pi/3 \\ 0 & x = \pi/3 \end{cases}$

3. $f(x) = \frac{|x|}{x}$

Examples

- (A) Which of the following have removable discontinuities? For those which do, what are the alternate functions with those discontinuities removed?
- (B) Which of the following have continuous extensions? For those which do, what are those extensions?

1. $f(x) = \frac{x^2 - 4}{x - 2}$ Cont. extension: $\bar{f}(x) = \begin{cases} f(x) & x \neq 2 \\ 4 & x = 2 \end{cases}$

2. $f(x) = \begin{cases} \sin x & x \neq \pi/3 \\ 0 & x = \pi/3 \end{cases}$ Removable disc.: $\bar{f}(x) = \sin(x)$

3. $f(x) = \frac{|x|}{x}$ No continuous extension.

One application: The Intermediate Value Theorem

Suppose f is continuous on a closed interval $[a, b]$.

If $f(a) < C < f(b)$ or $f(a) > C > f(b)$,

then there is at least one point c in the interval $[a, b]$ such that

$$f(c) = C.$$




One application: The Intermediate Value Theorem

Suppose f is continuous on a closed interval $[a, b]$.

If $f(a) < C < f(b)$ or $f(a) > C > f(b)$,

then there is at least one point c in the interval $[a, b]$ such that

$$f(c) = C.$$






One application: The Intermediate Value Theorem

Suppose f is continuous on a closed interval $[a, b]$.

If $f(a) < C < f(b)$ or $f(a) > C > f(b)$,

then there is at least one point c in the interval $[a, b]$ such that

$$f(c) = C.$$



Example 1: Show that the equation $x^5 - 3x + 1 = 0$ has at least one solution in the interval $[0, 1]$.

Example 2: Show every polynomial

$$p(x) = a_n x^n + \cdots + a_1 x + a_0, \quad a_n \neq 0$$

of odd degree has at least one real root (a solution to $p(x) = 0$).

Our favorite application: Rates of change!

It only makes sense to study the rate of change of a function where that function is continuous (or maybe where the function has a continuous extension)!

