



# The Definite Integral

## The Area Problem




## Upper and Lower Sums

Suppose we want to use rectangles to approximate the area under the graph of  $y = x + 1$  on the interval  $[0, 1]$ .



*Upper Riemann Sum*



*Lower Riemann Sum*

$$\frac{31}{20} > 1.5 > \frac{29}{20}$$

As you take more and more smaller and smaller rectangles, if  $f$  is nice, both of these will approach the real area.

$n$	$U$	$L$
100	1.505000000	1.495000000
150	1.503333333	1.496666667
200	1.502500000	1.497500000
300	1.501666667	1.498333333
500	1.501000000	1.499000000

## In general: finding the Area Under a Curve

Let  $y = f(x)$  be given and defined on an interval  $[a, b]$ .



## In general: finding the Area Under a Curve


Let  $y = f(x)$  be given and defined on an interval  $[a, b]$ .



Break the interval into  $n$  equal pieces.

## In general: finding the Area Under a Curve

Let  $y = f(x)$  be given and defined on an interval  $[a, b]$ .




Break the interval into  $n$  equal pieces.

Label the endpoints of those pieces  $x_0, x_1, \dots, x_n$ .

## In general: finding the Area Under a Curve

Let  $y = f(x)$  be given and defined on an interval  $[a, b]$ .



Break the interval into  $n$  equal pieces.


Label the endpoints of those pieces  $x_0, x_1, \dots, x_n$ .

Let  $\Delta x = x_i - x_{i-1} = \frac{b-a}{n}$  be the width of each interval.



## In general: finding the Area Under a Curve

Let  $y = f(x)$  be given and defined on an interval  $[a, b]$ .



Break the interval into  $n$  equal pieces.

Label the endpoints of those pieces  $x_0, x_1, \dots, x_n$ .


Let  $\Delta x = x_i - x_{i-1} = \frac{b-a}{n}$  be the width of each interval.

The **Upper Riemann Sum** is: let  $M_i$  be the *maximum* value of the function on that  $i^{\text{th}}$  interval, so

$$U(f, P) = M_1\Delta x + M_2\Delta x + \cdots + M_n\Delta x.$$

## In general: finding the Area Under a Curve

Let  $y = f(x)$  be given and defined on an interval  $[a, b]$ .



Break the interval into  $n$  equal pieces.

Label the endpoints of those pieces  $x_0, x_1, \dots, x_n$ .

Let  $\Delta x = x_i - x_{i-1} = \frac{b-a}{n}$  be the width of each interval.

The **Upper Riemann Sum** is: let  $M_i$  be the *maximum* value of the function on that  $i^{\text{th}}$  interval, so


$$U(f, P) = M_1 \Delta x + M_2 \Delta x + \cdots + M_n \Delta x.$$

The **Lower Riemann Sum** is: let  $m_i$  be the *minimum* value of the function on that  $i^{\text{th}}$  interval, so

$$L(f, P) = m_1 \Delta x + m_2 \Delta x + \cdots + m_n \Delta x.$$

## In general: finding the Area Under a Curve

Let  $y = f(x)$  be given and defined on an interval  $[a, b]$ .



Break the interval into  $n$  equal pieces.

Label the endpoints of those pieces  $x_0, x_1, \dots, x_n$ .

Let  $\Delta x = x_i - x_{i-1} = \frac{b-a}{n}$  be the width of each interval.


The **Upper Riemann Sum** is: let  $M_i$  be the *maximum* value of the function on that  $i^{\text{th}}$  interval, so

$$U(f, P) = M_1 \Delta x + M_2 \Delta x + \cdots + M_n \Delta x.$$


The **Lower Riemann Sum** is: let  $m_i$  be the *minimum* value of the function on that  $i^{\text{th}}$  interval, so

$$L(f, P) = m_1 \Delta x + m_2 \Delta x + \cdots + m_n \Delta x).$$

Take the limit as  $n \rightarrow \infty$  or  $\Delta x \rightarrow 0$ .



Upper



Lower

## Sigma Notation

If  $m$  and  $n$  are integers with  $m \leq n$ , and if  $f$  is a function defined on the integers from  $m$  to  $n$ , then the symbol  $\sum_{i=m}^n f(i)$ , called sigma notation, is means

$$\sum_{i=m}^n f(i) = f(m) + f(m+1) + f(m+2) + \cdots + f(n)$$

## Sigma Notation

If  $m$  and  $n$  are integers with  $m \leq n$ , and if  $f$  is a function defined on the integers from  $m$  to  $n$ , then the symbol  $\sum_{i=m}^n f(i)$ , called sigma notation, is means

$$\sum_{i=m}^n f(i) = f(m) + f(m+1) + f(m+2) + \cdots + f(n)$$

Examples:  $\sum_{i=1}^n i = 1 + 2 + 3 + \cdots + n$

$$\sum_{i=1}^n i^2 = 1^2 + 2^2 + 3^2 + \cdots + n^2$$

$$\sum_{i=1}^n \sin(i) = \sin(1) + \sin(2) + \sin(3) + \cdots + \sin(n)$$

$$\sum_{i=0}^{n-1} x^i = x^0 + x + x^2 + x^2 + x^3 + x^4 + \cdots + x^{n-1}$$

## Sigma Notation

If  $m$  and  $n$  are integers with  $m \leq n$ , and if  $f$  is a function defined on the integers from  $m$  to  $n$ , then the symbol  $\sum_{i=m}^n f(i)$ , called sigma notation, is means

$$\sum_{i=m}^n f(i) = f(m) + f(m+1) + f(m+2) + \cdots + f(n)$$

Examples:  $\sum_{i=1}^n i = 1 + 2 + 3 + \cdots + n$

$$\sum_{i=1}^n i^2 = 1^2 + 2^2 + 3^2 + \cdots + n^2$$

$$\sum_{i=1}^n \sin(i) = \sin(1) + \sin(2) + \sin(3) + \cdots + \sin(n)$$


$$\sum_{i=0}^{n-1} x^i = 1 + x + x^2 + x^2 + x^3 + x^4 + \cdots + x^{n-1}$$

# The Area Problem Revisited


$$\text{Upper Riemann Sum} = \sum_{i=1}^n M_i \Delta x$$

$$\text{Lower Riemann Sum} = \sum_{i=1}^n m_i \Delta x,$$

where  $M_i$  and  $m_i$  are, respectively, the maximum and minimum values of  $f$  on the  $i$ th subinterval  $[x_{i-1}, x_i]$ ,  $1 \leq i \leq n$ .



Upper





Lower

$$n = 5$$



## Example

Use an Upper Riemann Sum and a Lower Riemann Sum, first with 8, then with 100 subintervals of equal length to approximate the area under the graph of  $y = f(x) = x^2$  on the interval  $[0, 1]$ .



# The Definite Integral

We say that  $f$  is integrable on  $[a, b]$  if there exists a number  $A$  such that

$$\text{Lower Riemann Sum} \leq A \leq \text{Upper Riemann Sum}$$

for all  $n$ . We write the number as

$$A = \int_a^b f(x) dx$$

and call it the **definite integral** of  $f$  over  $[a, b]$ .

# The Definite Integral

We say that  $f$  is integrable on  $[a, b]$  if there exists a number  $A$  such that

$$\text{Lower Riemann Sum} \leq A \leq \text{Upper Riemann Sum}$$

for all  $n$ . We write the number as

$$A = \int_a^b f(x) dx$$

and call it the **definite integral** of  $f$  over  $[a, b]$ .

**Trickiness:** Who wants to find maxima/minima over every interval? Especially as  $n \rightarrow \infty$ ? Calculus nightmare!!

## More Riemann Sums

Let  $f$  be defined on  $[a, b]$ , and pick a positive integer  $n$ .

Let

$$\Delta x = \frac{b - a}{n}$$


## More Riemann Sums

Let  $f$  be defined on  $[a, b]$ , and pick a positive integer  $n$ .

Let

$$\Delta x = \frac{b - a}{n}$$

Notice:



$$x + 0 = a, \quad x_1 = a + \Delta x, \quad x_2 = a + 2\Delta x, \quad x_3 = a + 3\Delta x, \dots$$


## More Riemann Sums

Let  $f$  be defined on  $[a, b]$ , and pick a positive integer  $n$ .

Let

$$\Delta x = \frac{b - a}{n}$$

Notice:



$$x + 0 = a, \quad x_1 = a + \Delta x, \quad x_2 = a + 2\Delta x, \quad x_3 = a + 3\Delta x, \dots$$

So let

$$x_i = a + i * \Delta x.$$

## More Riemann Sums

Let  $f$  be defined on  $[a, b]$ , and pick a positive integer  $n$ .

Let

$$\Delta x = \frac{b - a}{n} \quad \text{and} \quad x_i = a + i * \Delta x.$$

## More Riemann Sums


Let  $f$  be defined on  $[a, b]$ , and pick a positive integer  $n$ .

Let

$$\Delta x = \frac{b - a}{n} \quad \text{and} \quad x_i = a + i * \Delta x.$$


Then the Right Riemann Sum is

$$\sum_{i=1}^n f(x_i) \Delta x,$$



and the Left Riemann Sum is

$$\sum_{i=0}^{n-1} f(x_i) \Delta x.$$





# Integrals made easier

## **Theorem**

*If  $f$  is continuous on  $[a, b]$ , then  $f$  is Riemann integrable on  $[a, b]$ .*

# Integrals made easier

## Theorem

*If  $f$  is continuous on  $[a, b]$ , then  $f$  is Riemann integrable on  $[a, b]$ .*

## Theorem

*If  $f$  is Riemann integrable on  $[a, b]$ , then*

$$\int_a^b f(x) dx = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(c_i) \Delta x_i$$

*where  $c_i$  is **any** point in the interval  $[x_{i-1}, x_i]$ .*

# Integrals made easier

## Theorem

If  $f$  is continuous on  $[a, b]$ , then  $f$  is Riemann integrable on  $[a, b]$ .

## Theorem

If  $f$  is Riemann integrable on  $[a, b]$ , then

$$\int_a^b f(x) dx = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(c_i) \Delta x_i$$

where  $c_i$  is **any** point in the interval  $[x_{i-1}, x_i]$ .

**Punchline:** (1) Every continuous function has an integral, and (2) we can get there by just using right or left sums! (instead of upper or lower sums)

# Properties of the Definite Integral

1.  $\int_a^a f(x)dx = 0.$


2. If  $f$  is integrable and

(a)  $f(x) \geq 0$  on  $[a, b]$ , then  $\int_a^b f(x)dx$  equals the area of the region under the graph of  $f$  and above the interval  $[a, b]$ ;

(b)  $f(x) \leq 0$  on  $[a, b]$ , then  $\int_a^b f(x)dx$  equals the **negative** of the area of the region between the interval  $[a, b]$  and the graph of  $f$ .


3.  $\int_b^a f(x)dx = - \int_a^b f(x)dx.$

4. If  $a < b < c$ ,  $\int_a^b f(x)dx + \int_b^c f(x)dx = \int_a^c f(x)dx$



5. If  $f$  is an **even** function, then


$$\int_{-a}^a f(x) dx = 2 \int_0^a f(x) dx.$$



Area I = Area II

6. If  $f$  is an **odd** function, then


$$\int_{-a}^a f(x) dx = 0.$$



Area I = Area II

## Example

$$\text{If } f(x) = \begin{cases} x, & x < 0, \\ \sqrt{1 - (x - 1)^2}, & 0 \leq x \leq 2, \\ x - 2, & x \geq 2, \end{cases} \text{ what is } \int_{-1}^3 f(x) dx?$$





# Mean Value Theorem for Definite Integrals

## Theorem

Let  $f$  be continuous on the interval  $[a, b]$ . Then there exists  $c$  in  $[a, b]$  such that

$$\int_a^b f(x)dx = (b - a)f(c).$$

## Definition

The *average value* of a continuous function on the interval  $[a, b]$  is

$$\frac{1}{b - a} \int_a^b f(x)dx.$$