

Modeling with Differential Equations: Introduction to the Issues

Warm-up

Do you know a function...

... whose first derivative is the same as the function itself, i.e.

$$\frac{d}{dx}f(x) = f(x)?$$

... whose first derivative is negative of the function, i.e.

$$\frac{d}{dx}f(x) = -f(x)?$$

... whose second derivative is negative of itself, i.e.

$$\frac{d^2}{dx^2}f(x) = -f(x)?$$

Goal:

Given an equation relating a variable (e.g. x), a function (e.g. y), and its derivatives (y', y'', \dots), **what is y ?**
i.e. How do I solve for y ?

Goal:

Given an equation relating a variable (e.g. x), a function (e.g. y), and its derivatives (y' , y'' , \dots), **what is y ?**
i.e. How do I solve for y ?

Why?

Many physical and biological systems can be modeled with differential equations. Also, it can be a lot harder to model a function long term than it is to measure how something changes as the system goes from one state to another.

Some examples

Obvervation: The rate of increase of a bacterial culture is proportional to the number of bacteria present at that time.

Obvervation: The motion of a mass on a spring is given by two opposing forces: (1) the force exerted by the mass in motion ($F = ma = m \frac{d^2}{dt^2} D$) and (2) the force exerted by the spring, proportional to the displacement from equilibrium ($F = kD$).

Some examples

Obvervation: The rate of increase of a bacterial culture is proportional to the number of bacteria present at that time.

Equation: $\frac{dP}{dt} = kP$

Obvervation: The motion of a mass on a spring is given by two opposing forces: (1) the force exerted by the mass in motion ($F = ma = m \frac{d^2}{dt^2} D$) and (2) the force exerted by the spring, proportional to the displacement from equilibrium ($F = kD$).

Equation: $m * \frac{d^2}{dt^2} D = -kD$

Some examples

Obvervation: The rate of increase of a bacterial culture is proportional to the number of bacteria present at that time.

Equation: $\frac{dP}{dt} = kP$

Solution: $P = Ae^{kt}$, where A is a constant.

Obvervation: The motion of a mass on a spring is given by two opposing forces: (1) the force exerted by the mass in motion ($F = ma = m \frac{d^2}{dt^2} D$) and (2) the force exerted by the spring, proportional to the displacement from equilibrium ($F = kD$).

Equation: $m * \frac{d^2}{dt^2} D = -kD$

Solution: $D(t) = A \cos(t * \sqrt{k/m}) + B \sin(t * \sqrt{k/m})$,
where A , B , k , and m are all constants.

Slope Fields

If you can write your differential equation like

$$\frac{dy}{dx} = F(x, y)$$

then you really have a way of saying

“If I’m standing at the point (a, b) ,
then I should move from here with slope $F(a, b)$.”

Slope Fields

If you can write your differential equation like

$$\frac{dy}{dx} = F(x, y)$$

then you really have a way of saying

“If I’m standing at the point (a, b) ,
then I should move from here with slope $F(a, b)$.”

Some examples:

$$\frac{dy}{dx} = -\frac{x}{y}$$

$$\frac{dP}{dt} = kP$$

$$\frac{dx}{dt} = t^2 \sin(xt) + x^2$$

Slope Fields

If you can write your differential equation like

$$\frac{dy}{dx} = F(x, y)$$

then you really have a way of saying

“If I’m standing at the point (a, b) ,
then I should move from here with slope $F(a, b)$.”

Some examples:

$$\frac{dy}{dx} = -\frac{x}{y}$$

$$\frac{dP}{dt} = kP$$

$$\frac{dx}{dt} = t^2 \sin(xt) + x^2$$

Some non-examples:


$$\frac{dy}{dx} = -\frac{x}{y} + \frac{d^2y}{dx^2}$$

$$\frac{dP}{dt} * \frac{d^2P}{dt^2} = kP$$

$$m * \frac{d^2D}{dt^2} = -kD$$


x	y	$\frac{dy}{dx} = -x/y$
0	1	
0	-1	
1	1	
1	-1	
-1	1	
-1	-1	
2	1	
1	2	
-2	0	

Slope field:




x	y	$\frac{dy}{dx} = -x/y$
0	1	0
0	-1	
1	1	
1	-1	
-1	1	
-1	-1	
2	1	
1	2	
-2	0	

Slope field:




x	y	$\frac{dy}{dx} = -x/y$
0	1	0
0	-1	0
1	1	
1	-1	
-1	1	
-1	-1	
2	1	
1	2	
-2	0	

Slope field:




x	y	$\frac{dy}{dx} = -x/y$
0	1	0
0	-1	0
1	1	-1
1	-1	
-1	1	
-1	-1	
2	1	
1	2	
-2	0	

Slope field:




x	y	$\frac{dy}{dx} = -x/y$
0	1	0
0	-1	0
1	1	-1
1	-1	1
-1	1	
-1	-1	
2	1	
1	2	
-2	0	


Slope field:




x	y	$\frac{dy}{dx} = -x/y$
0	1	0
0	-1	0
1	1	-1
1	-1	1
-1	1	1
-1	-1	-1
2	1	-2
1	2	-1/2
-2	0	undef




x	y	$\frac{dy}{dx} = -x/y$
0	1	0
0	-1	0
1	1	-1
1	-1	1
-1	1	1
-1	-1	-1
2	1	-2
1	2	-1/2
-2	0	undef



x	y	$\frac{dy}{dx} = -x/y$
0	1	0
0	-1	0
1	1	-1
1	-1	1
-1	1	1
-1	-1	-1
2	1	-2
1	2	-1/2
-2	0	undef




x	y	$\frac{dy}{dx} = -x/y$
0	1	0
0	-1	0
1	1	-1
1	-1	1
-1	1	1
-1	-1	-1
2	1	-2
1	2	-1/2
-2	0	undef



Arrows point in the direction of semicircles! $y = \pm\sqrt{r^2 - x^2}$?


x	y	$\frac{dy}{dx} = -x/y$
0	1	0
0	-1	0
1	1	-1
1	-1	1
-1	1	1
-1	-1	-1
2	1	-2
1	2	-1/2
-2	0	undef



Arrows point in the direction of semicircles! $y = \pm\sqrt{r^2 - x^2}$?

Check:
$$\frac{d}{dx} \pm \sqrt{r^2 - x^2} = \frac{-2x}{\pm 2\sqrt{r^2 - x^2}}$$

x	y	$\frac{dy}{dx} = -x/y$
0	1	0
0	-1	0
1	1	-1
1	-1	1
-1	1	1
-1	-1	-1
2	1	-2
1	2	-1/2
-2	0	undef



Arrows point in the direction of semicircles! $y = \pm\sqrt{r^2 - x^2}$?

Check: $\frac{d}{dx} \pm \sqrt{r^2 - x^2} = \frac{-2x}{\pm 2\sqrt{r^2 - x^2}} = -\frac{x}{y} \quad \text{☺}$

Solving explicitly (get a formula!)

We've done...

1. Get lucky

“what's a function you know whose derivative blah blah ...”

2. Differential equations of the form

$$\frac{dy}{dx} = f(x)$$

Find the antiderivative!

Today, we'll add

3. Differential equations of the form

$$\frac{dy}{dx} = f(x) * g(y)$$

Use **“Separation of Variables”**

Separable Equations

A **separable differential equation** is one of the form

$$\frac{dy}{dx} = f(x) \cdot g(y).$$

Separable Equations

A **separable differential equation** is one of the form

$$\frac{dy}{dx} = f(x) \cdot g(y).$$

Some examples:

$$\frac{dy}{dx} = -\frac{x}{y} = (-x) * \left(\frac{1}{y}\right)$$

$$\frac{dx}{dt} = t^2 \sec(x)$$

Separable Equations

A **separable differential equation** is one of the form

$$\frac{dy}{dx} = f(x) \cdot g(y).$$

Some examples:

$$\frac{dy}{dx} = -\frac{x}{y} = (-x) * \left(\frac{1}{y}\right)$$

$$\frac{dx}{dt} = t^2 \sec(x)$$

Some non-examples:

$$\frac{dy}{dx} = x + y$$

$$\frac{dx}{dt} = \frac{t + x}{xt^2}$$

Separable Equations

A **separable differential equation** is one of the form

$$\frac{dy}{dx} = f(x) \cdot g(y).$$

Some examples:

$$\frac{dy}{dx} = -\frac{x}{y} = (-x) * \left(\frac{1}{y}\right)$$

$$\frac{dx}{dt} = t^2 \sec(x)$$

Some non-examples:

$$\frac{dy}{dx} = x + y$$

$$\frac{dx}{dt} = \frac{t + x}{xt^2}$$

A separable equation is one in which we can put all of the y 's and dy 's (as products) on one side of the equation and all of the x 's and dx 's (as products) on the other...

Examples

(1) If $\frac{dy}{dx} = -\frac{x}{y}$, then $y \, dy = -x \, dx$.

Examples

(1) If $\frac{dy}{dx} = -\frac{x}{y}$, then $y \, dy = -x \, dx$.

(2) If $\frac{dx}{dt} = t^2 \sec(x)$, then $\cos(x) \, dx = t^2 \, dt$.

Examples

$$(1) \text{ If } \frac{dy}{dx} = -\frac{x}{y}, \quad \text{then } y \, dy = -x \, dx.$$

$$(2) \text{ If } \frac{dx}{dt} = t^2 \sec(x), \quad \text{then } \cos(x) \, dx = t^2 \, dt.$$

To solve (1), integrate both sides:

$$\int y \, dy = \int -x \, dx$$

Examples

$$(1) \text{ If } \frac{dy}{dx} = -\frac{x}{y}, \quad \text{then } y \, dy = -x \, dx.$$

$$(2) \text{ If } \frac{dx}{dt} = t^2 \sec(x), \quad \text{then } \cos(x) \, dx = t^2 \, dt.$$

To solve (1), integrate both sides:

$$y^2/2 + c_2 = \int y \, dy = \int -x \, dx = -x^2/2 + c_1$$

Examples

$$(1) \text{ If } \frac{dy}{dx} = -\frac{x}{y}, \quad \text{then } y \, dy = -x \, dx.$$

$$(2) \text{ If } \frac{dx}{dt} = t^2 \sec(x), \quad \text{then } \cos(x) \, dx = t^2 \, dt.$$

To solve (1), integrate both sides:

$$y^2/2 + c_2 = \int y \, dy = \int -x \, dx = -x^2/2 + c_1$$


So

$$y = \pm \sqrt{2(-x^2/2 + c_1 - c_2)} = \pm \sqrt{a - x^2}$$

where $a = 2(c_1 - c_2)$.

Examples

Slope field for $\frac{dy}{dx} = -\frac{x}{y}$:



Suggested and checked $y = \pm\sqrt{r^2 - x^2}$

Examples

$$(1) \text{ If } \frac{dy}{dx} = -\frac{x}{y}, \quad \text{then } y \, dy = -x \, dx.$$

$$(2) \text{ If } \frac{dx}{dt} = t^2 \sec(x), \quad \text{then } \cos(x) \, dx = t^2 \, dt.$$

To solve (1), integrate both sides:

$$y^2/2 + c_2 = \int y \, dy = \int -x \, dx = -x^2/2 + c_1$$

So

$$y = \pm \sqrt{2(-x^2/2 + c_1 - c_2)} = \pm \sqrt{a - x^2}$$

where $a = 2(c_1 - c_2)$.

Find an implicit formula for (2) (with no derivatives left in it)

How many solutions are there?

Existence?

How do I know I even get a solution?

An important result in the theory of differential equations is

Peano's Existence Theorem, which states. . .

If $\frac{dy}{dx} = F(x, y)$ and $y(a) = b$,
where $F(x, y)$ is continuous in a domain D ,
then there is always **at least one solution** in the
domain, and any such solution is differentiable.

Uniqueness?

How do we know that there is not another solution?


If, additionally, $F(x, y) = f(x)g(y)$, and if g'
and f' are continuous, then solution is unique.

Example of non-uniqueness

Suppose $\frac{dy}{dx} = \frac{1}{x}$ and $y(2) = 1$.


Example of non-uniqueness

Suppose $\frac{dy}{dx} = \frac{1}{x}$ and $y(2) = 1$.




Example of non-uniqueness

Suppose $\frac{dy}{dx} = \frac{1}{x}$ and $y(2) = 1$.




Example of non-uniqueness

Suppose $\frac{dy}{dx} = \frac{1}{x}$ and $y(2) = 1$.




Example of non-uniqueness

Suppose $\frac{dy}{dx} = \frac{1}{x}$ and $y(2) = 1$.




Example of non-uniqueness

Suppose $\frac{dy}{dx} = \frac{1}{x}$ and $y(2) = 1$.




Example of non-uniqueness

Suppose $\frac{dy}{dx} = \frac{1}{x}$ and $y(2) = 1$.



Example of non-uniqueness





Suppose $\frac{dy}{dx} = \frac{1}{x}$ and $y(2) = 1$.



Since $\frac{1}{x}$ is not continuous at $x = 0$, we might have lots of solutions, all that split at 0!

(1) Match the differential equations to the slope fields:

(A) $\frac{dy}{dx} = \frac{1}{5}xy$ (B) $\frac{dy}{dx} = x+y$ (C) $\frac{dy}{dx} = \cos(x)$ (D) $\frac{dy}{dx} = \cos(y)$



(2) Solve the initial value problems

(a) $\frac{dy}{dx} = \frac{1}{5}xy, \quad y(0) = 2;$

(b) $\frac{dy}{dx} = \sin(x)/y^2, \quad y(0) = 3.$