

MATH 56 WORKSHEET : Euler-Machin formulae for  $\pi$ 

Start w/ addition formula for tangent:  $\tan(x+y) = \frac{\tan x + \tan y}{1 - \tan x \tan y}$

A) Set  $\alpha = \tan x$  &  $\tan^{-1}$  both sides:  
 $\beta = \tan y$

B) Check  $\alpha = 1/2$ ,  $\beta = 1/3$  works (Euler)

If want to push  $\alpha \rightarrow 0$ , what happens to  $\beta$  if result is to be  $\tan^{-1} 1$ ?

C) We can further split  $\tan^{-1} 1/2 = \tan^{-1} 1/3 + \tan^{-1}(\text{something})$

Solve for "something":

Thus write  $\pi/4 = \dots$

Why is this preferred?

D) to get  $\pi$  to  $N$  digits how many Taylor terms needed? (big-O)

So, what is total complexity? (big-O)

# MATH 56 WORKSHEET : Euler-Machin Formulae for $\pi$

## SOLUTIONS

Start w/ addition formula for tangent:  $\tan(x+y) = \frac{\tan x + \tan y}{1 - \tan x \tan y}$

A) Set  $\alpha = \tan x$  &  $\tan^{-1}$  both sides:  $x = \tan^{-1} \alpha$   $y = \tan^{-1} \beta$   
 $\beta = \tan y$

$$\tan^{-1} \alpha + \tan^{-1} \beta = \frac{\alpha + \beta}{1 - \alpha\beta}$$

B) Check  $\alpha = 1/2$ ,  $\beta = 1/3$  works (Euler)  $\frac{\alpha + \beta}{1 - \alpha\beta} = \frac{1/2 + 1/3}{1 - 1/6} = \frac{5/6}{5/6} = 1$   
 so  $\tan^{-1} 1/2 + \tan^{-1} 1/3 = \tan^{-1} 1$ .

If want to push  $\alpha \rightarrow 0$ , what happens to  $\beta$  if result is to be  $\tan^{-1} 1$ ?

Fix  $\frac{\alpha + \beta}{1 - \alpha\beta} = 1$ , solve for  $\beta$ :  $\beta = \frac{1 - \alpha}{1 + \alpha}$  so as  $\alpha \rightarrow 0^+$ ,  $\beta \rightarrow 1^-$

C) We can further split  $\tan^{-1} 1/2 = \tan^{-1} 1/3 + \tan^{-1}(\text{something})$

Solve for "something":  $\frac{\alpha + \beta}{1 - \alpha\beta} = 1/2$  so  $\beta = \frac{1/2 - \alpha}{1 + \alpha/2} = \frac{1/2 - 1/3}{1 + 1/6} = \frac{1/6}{7/6} = 1/7$

Thus write  $\pi/4 = 2 \tan^{-1} 1/3 + \tan^{-1} 1/7 = 2 \left[ \frac{1}{13} - \frac{1}{3 \cdot 3^3} + \frac{1}{5 \cdot 3^5} - \dots \right] + \frac{1}{1 \cdot 7} - \frac{1}{3 \cdot 7^3} + \frac{1}{5 \cdot 7^5} - \dots$

Why is this preferred? convergence rate is largest  $|x| = 1/3 = r$ , ie error =  $O(1/3^n)$ , beats  $r = 1/2$  above

D) to get  $\pi$  to  $N$  digits, how many Taylor terms needed? (big-O)

So, what is total complexity? (big-O)  $O(N^2)$ , assuming  $O(n)$  to update  $\frac{1}{3^n} \rightarrow \frac{1}{3^{n+1}}$  each term.  
 $\hookrightarrow \left(\frac{1}{3}\right)^2 \approx \frac{1}{10}$  so roughly  $2N = O(N)$  terms.